

University of Dundee

AMP-Activated Protein Kinase

Salt, Ian P.; Hardie, D. Grahame

Published in:
Circulation Research

DOI:
[10.1161/CIRCRESAHA.117.309633](https://doi.org/10.1161/CIRCRESAHA.117.309633)

Publication date:
2017

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Salt, I. P., & Hardie, D. G. (2017). AMP-Activated Protein Kinase: An Ubiquitous Signaling Pathway With Key Roles in the Cardiovascular System. *Circulation Research*, 120(11), 1825-1841.
<https://doi.org/10.1161/CIRCRESAHA.117.309633>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

AMP-ACTIVATED PROTEIN KINASE – A UBIQUITOUS SIGNALLING PATHWAY WITH KEY ROLES IN THE CARDIOVASCULAR SYSTEM

Ian P. Salt¹ and D. Grahame Hardie²

¹Institute of Cardiovascular & Medical Sciences, College of Medical, Veterinary & Life Sciences,
Davidson Building, University of Glasgow, Glasgow, G12 8QQ, Scotland, UK

²Division of Cell Signalling & Immunology, School of Life Sciences, University of Dundee,
Dundee, DD1 5EH, Scotland, UK

Short title: AMPK: roles in cardiovascular system

KEYWORDS: AMPK, AMP-activated protein kinase, heart, heart defects (congenital),
metabolism, vasculature

ABSTRACT

The AMP-activated protein kinase (AMPK) is a key regulator of cellular and whole body energy homeostasis, which acts to restore energy homeostasis whenever cellular energy charge is depleted. Over the last two decades, it has become apparent that AMPK regulates a number of other cellular functions and has specific roles in cardiovascular tissues, acting to regulate cardiac metabolism and contractile function as well as promoting anti-contractile, anti-inflammatory and anti-atherogenic actions in blood vessels. In this review, we will discuss the role of AMPK in the cardiovascular system, including the molecular basis of mutations in AMPK that alter cardiac physiology and the proposed mechanisms by which AMPK regulates vascular function under physiological and pathophysiological conditions.

The AMP-activated protein kinase (AMPK) is the central component of a signalling pathway that regulates the switch between anabolism and catabolism, as well as many other aspects of cell function¹⁻³. In this review we focus on the physiological roles of AMPK within the cardiovascular system, although we will start by discussing general features that apply in all mammalian cell types. Readers are also referred to two other recent reviews of the role of AMPK in cardiovascular disease^{4, 5}.

STRUCTURE AND REGULATION OF AMPK

Occurrence of subunit isoforms

AMPK exists universally as heterotrimeric complexes containing catalytic α subunits and regulatory β and γ subunits, which occur as multiple isoforms ($\alpha 1/\alpha 2$; $\beta 1/\beta 2$; $\gamma 1/\gamma 2/\gamma 3$) encoded by distinct genes. These could give rise in principle to twelve heterotrimeric combinations, although specific combinations appear to be favoured in specific cell types. For example, although skeletal muscle expresses mRNAs encoding all seven subunit isoforms, assays of immunoprecipitated isoforms suggest that AMPK activity in human skeletal muscle is accounted for by just three combinations: $\alpha 1\beta 2\gamma 1$, $\alpha 2\beta 2\gamma 1$ and $\alpha 2\beta 2\gamma 3$ ⁶.

The domain organization of the seven AMPK subunit isoforms are shown in Fig. 1, and a representation of a crystal structure for the human $\alpha 1\beta 2\gamma 1$ heterotrimer⁷ is shown in Fig. 2. Similar structures of the $\alpha 2\beta 1\gamma 1$ ⁸ and $\alpha 1\beta 1\gamma 1$ ⁹ complexes are available.

The α subunits

The α subunits contain a kinase domain (α -KD) at their N-termini, with the small N-lobe (yellow in Figs.1 and 2) and larger C-lobe (green) typical of all protein kinases. The substrate Mg.ATP^{2-} binds in a deep cleft between these lobes, occupied in the structure in Fig. 2 by the non-specific kinase inhibitor staurosporine. Isolated α -KDs are only significantly active after phosphorylation by upstream kinases at a conserved threonine, usually referred to as Thr172. Thr172 phosphorylation is a good marker for AMPK activity, and Western blotting using phosphospecific antibodies against this site is often used as a semi-quantitative measure of AMPK activity. However, this ignores the effects of allosteric activation of AMPK, a problem that can be overcome by also addressing the phosphorylation state of a downstream target such as acetyl-CoA carboxylase (ACC).

The α -KD is followed by a small globular domain termed the autoinhibitory domain (AID, orange) that, when AMP is not bound to the γ subunit, binds to both the N- and C-lobes of the KD and thus causes inhibition⁷. The AID is connected to the C-terminal domain (α -CTD, red) by the α -linker (blue), a region of extended polypeptide that interacts with the γ subunit when AMP is bound at site 3 (as is the case in Fig. 2), thus pulling the AID away from this inhibitory position.

The β subunits

Two well-conserved regions within the β subunits are the C-terminal domain (β -CTD, silver-grey), which forms the “core” of the heterotrimeric complex, and the central carbohydrate-binding module (β -CBM, silver-blue). The β -CBM is related to carbohydrate-binding domains usually occurring in enzymes that metabolize starch or glycogen, which localize those enzymes on their polysaccharide substrate. Consistent with this, the β -CBM causes a proportion of AMPK in cells to bind to the surface of glycogen particles¹⁰⁻¹². In all three structures of mammalian heterotrimers, the β -CBM interacts with the N-lobe of the α subunit KD, with its glycogen-binding site (defined by binding of the oligosaccharide β -cyclodextrin) at the top in the view of Fig. 2.

What is the function of glycogen binding by AMPK? The muscle and liver isoforms of glycogen synthase, which are also bound to glycogen particles, are physiological targets for AMPK^{13, 14}, and one function may be to co-localize AMPK with them. The β -CBM is also of interest because the cleft between it and the N-lobe of the α subunit forms the binding site for activators such as A769662 and 991⁸. This site has been termed the Allosteric Drug and Metabolite (ADaM) site¹⁵, and is discussed further below.

The γ subunits

All γ subunits contain four tandem repeats of a sequence motif known as a CBS motif. In other proteins, pairs of CBS motifs often form binding sites for regulatory ligands containing adenosine¹⁶; in the AMPK- γ subunits they bind the regulatory nucleotides AMP, ADP or and ATP¹⁷. Two views of a structure for the γ 1 subunit, containing three molecules of bound AMP¹⁸, are shown in Fig. 3. The two pairs of repeats (CBS1:CBS2 and CBS3:CBS4) assemble in a pseudosymmetrical “head-to-head” manner. The γ subunit thus forms a structure like a flattened disc (seen from different faces in Fig. 3) with one CBS repeat in each quadrant, generating four potential ligand-binding clefts in the centre. However, one of these appears to be unused, perhaps because conserved

aspartate residues in CBS1, CBS3 and CBS4 that bind the ribose ring of adenine nucleotides in sites 1, 3 and 4 are absent from CBS2¹⁸. Interestingly, mutation of any of these three aspartate residues interferes with multiple effects of AMP on kinase activity¹⁹, suggesting that occupancy of all three sites may be required for full activation.

Identity of upstream kinases

Identifying the upstream kinases that phosphorylate Thr172 was a difficult challenge, eventually solved by genome-wide biochemical screens that identified three upstream kinases in budding yeast^{20, 21}. The mammalian kinases with catalytic domains most closely related to these were LKB1 and the Ca²⁺/calmodulin-dependent kinase CaMKK2 (CaMKK β), and evidence was soon obtained that both could act as physiological upstream kinases in mammalian cells²²⁻²⁷. The discovery that LKB1 was an upstream kinase for AMPK was particularly interesting, because LKB1 had previously been identified to be a tumor suppressor²⁸. Phosphorylation of Thr172 by CaMKK2²⁵⁻²⁷ represents a mechanism by which hormones that increase cytosolic Ca²⁺ can activate AMPK in the absence of energy stress.

GENERAL FEATURES OF AMPK REGULATION

Regulation by the canonical energy stress mechanism

It might be expected that a system that monitors cellular energy status would sense ATP and ADP but, interestingly, all metabolic enzymes known to directly monitor cellular energy charge (glycogen phosphorylase, 6-phosphofructo-1-kinase, fructose-1,6-bisphosphatase) primarily sense AMP and ATP, as does AMPK. The principal source of AMP in cells is thought to be the adenylate kinase reaction ($2\text{ADP} \leftrightarrow \text{ATP} + \text{AMP}$), which appears to operate close to equilibrium in most cells so that the AMP:ATP ratio will vary as the square of the ADP:ATP ratio²⁹. The former is therefore a more sensitive indicator of falling energy status than the latter.

Binding of AMP activates AMPK by three complementary mechanisms, of which the first two are mimicked by ADP at higher concentration, while all three are antagonized by ATP: (i) inhibition of Thr172 dephosphorylation by protein phosphatases; (ii) promotion of Thr172 phosphorylation by LKB1; and (iii) allosteric activation^{30, 31}. The structural model shown in Fig. 2 suggests a mechanism, for which there is now supporting evidence⁷, to explain mechanisms (i) and

(iii). When AMP is bound at site 3, the α -linker interacts with the surface of the γ subunit containing that site (Fig. 2)^{7, 32}. AMPK heterotrimers contain two rather distinct regions, the “catalytic module” (β -CBM, KD, AID, top/front section in Fig. 2) and the “nucleotide-binding module” (γ subunit, β -CTD, α -CTD, bottom/rear section in Fig. 2). The “hinge” connecting them is the α -linker, and release of the α -linker on binding of ATP rather than AMP at site 3 is envisaged to allow the two modules to move apart, causing the AID to rotate back into its inhibitory position behind the α -KD. This conformational change would also increase accessibility of Thr172 to protein phosphatases, which in the AMP-bound conformation of Fig. 2 is around the back, located in a deep cleft between the two modules. This model therefore explains not only how AMP binding at site 3 causes allosteric activation, but also why it protects against Thr172 dephosphorylation, with binding of ATP antagonizing both effects.

Multiple mechanisms of pharmacological activation of AMPK

A selection of compounds commonly used to activate AMPK experimentally are listed in Table 1. Those in Class 1, including the antidiabetic drug metformin and berberine (derived from traditional Chinese medicine) inhibit Complex I of the mitochondrial respiratory chain, while those in class 2, including 2-deoxyglucose, inhibit glycolysis. Both classes activate AMPK indirectly by increasing cellular AMP:ATP ratios³³. These compounds are frequently used to activate AMPK experimentally, and studies utilizing them are described later in this review. However, because they work by depleting cellular ATP, they should not be regarded as specific AMPK activators and any results obtained with them should ideally be followed up using genetic approaches, such as the use of AMPK knockouts.

The third class of activator includes the widely used compound 5-aminoimidazole-4-carboxamide ribonucleoside (AICAR), a nucleoside that is taken up into cells and converted to the equivalent nucleotide, ZMP. An important caveat here is that although ZMP is an AMP analog that mimics the effects of AMP to activate AMPK, it is about 50-fold less potent than AMP itself³⁴. Because ZMP accumulates to millimolar concentrations inside cells AICAR does cause AMPK activation, but ZMP also has off-target effects. For example, it is known to mimic the effects of AMP on phosphorylase³⁵ and fructose-1,6-bisphosphatase³⁶ as well as AMPK. Another problem with AICAR is that it is an adenosine analogue and, while it does not appear to bind directly to

141 adenosine receptors, in incubated cell systems it competes with endogenous adenosine for reuptake
142 into cells by adenosine transporters, so can have adenosine-like effects³⁷.

143 A more specific activator that works via a related mechanism (class 4) is C13, which is taken up
144 by cells and converted by cellular esterases to the AMP analog C2, a very potent AMPK activator³⁸
145 although only for α 1- and not α 2-containing complexes³⁹. Despite its selectivity for the α 1 isoform,
146 C2 binds to the γ subunit. Although the AMP and C2 binding sites overlap, they are not identical⁴⁰.

147 The other molecules of choice for selective activation of AMPK are those binding at the ADaM
148 site located between the β -CBM and the N-lobe on the α subunit, such as A769662 and 991 (class
149 5)⁸. All compounds binding this site are more potent activators of complexes containing AMPK- β 1
150 rather than β 2, and most are essentially β 1-selective (Table 1). Using AMPK phosphorylated on
151 Thr172, these activators cause a modest degree of allosteric activation (up to 4-fold) and, by
152 inhibiting dephosphorylation, also promote net Thr172 phosphorylation^{41, 42}. More remarkably, they
153 cause a much larger allosteric activation (up to 65-fold) of AMPK that is not phosphorylated on
154 Thr172, and this effect is synergistic with AMP⁴³. At present, almost all of the activators known to
155 bind at this site are synthetic compounds derived from high-throughput screens, although there is
156 much speculation in the field that there is a naturally occurring ligand, hence the appearance of
157 “metabolite” in its name¹⁵. However, no natural ligands occurring in animal cells that bind the
158 ADaM site have yet been found. Salicylate, the natural plant product from which acetyl salicylic
159 acid (ASA, aspirin) was derived, does activate AMPK by binding this site^{9, 44}.

160 It should also be noted that there are currently no specific pharmacological inhibitors of AMPK;
161 compound C (dorsomorphin) is sometimes claimed to be a specific inhibitor, but this is not the
162 case⁴⁵.

163 **TARGETS AND PATHWAYS DOWNSTREAM OF AMPK**

164 ***Catabolic effects switched on by AMPK***

165 Once activated by energy stress, AMPK switches on catabolic pathways generating ATP, while
166 switching off anabolic pathways and other processes consuming ATP, thus acting to restore energy
167 homeostasis. This topic has been discussed in more detail in previous reviews¹⁻³. Examples of
168 catabolic processes acutely switched on include cellular glucose uptake mediated by GLUT1 and
169 GLUT4. Activation of GLUT1 may occur via phosphorylation of the thioredoxin interacting protein

170 TXNIP⁴⁶, while enhanced GLUT4 translocation to the plasma membrane in muscle appears to
171 occur, at least in part, by phosphorylation of TBC1D1, which modulates trafficking of GLUT4-
172 containing vesicles⁴⁷. AMPK activation also enhances GLUT4 expression⁴⁸, in part via
173 phosphorylation of the histone deacetylase HDAC5, promoting binding of 14-3-3 proteins and
174 consequent retention of HDAC5, a transcriptional inhibitor, in the cytoplasm^{49, 50}. AMPK can also
175 cause a short-term activation of glycolysis via phosphorylation of PFKFB2⁵¹ and PFKFB3⁵²,
176 isoforms of the bifunctional enzyme that synthesizes and breaks down fructose-2,6-bisphosphate, a
177 key allosteric activator of 6-phosphofructo-1-kinase and hence glycolysis. PFKFB2 is expressed in
178 the heart, while PFKFB3 occurs as an “inducible” form whose expression in monocytes and
179 macrophages is induced by inflammatory mediators such as lipopolysaccharide⁵³. Phosphorylation
180 of PFKFB2 or PFKFB3 at equivalent sites near their C-termini increases the synthesis of fructose-
181 2,6-bisphosphate and hence promotes glycolysis during hypoxia in heart and in activated
182 monocytes/macrophages, respectively. This may enhance survival of these cells during periods of
183 hypoxia or ischemia.

184 Although AMPK can therefore activate glucose uptake and glycolysis in specific cell types, in
185 the longer term it tends to promote instead the more glucose-sparing and energy-efficient oxidative
186 metabolism. In skeletal muscle^{54, 55} and liver⁵⁶, AMPK activates fatty acid oxidation by inhibiting
187 the ACC1/ACC2 isoforms of acetyl-CoA carboxylase to reduce malonyl-CoA, an inhibitor of the
188 uptake of fatty acids into mitochondria. AMPK promotes the expression of TCA cycle enzymes⁵⁷,
189 as well as mitochondrial biogenesis, which it achieves by increasing the expression/activity of the
190 transcriptional co-activator PGC-1 α , either by direct phosphorylation triggering a positive feedback
191 effect on its own expression⁵⁸, or by enhancing its deacetylation by SIRT1⁵⁹. PGC-1 α acts as a co-
192 activator for several transcription factors involved in mitochondrial biogenesis and oxidative
193 metabolism, including myocyte enhancer factor-2 (MEF2), nuclear respiratory factors-1/-2
194 (NRF1/2), and PPAR- α and - δ ⁶⁰. Despite this evidence that AMPK promotes oxidative
195 metabolism, in mice with a skeletal/cardiac muscle-specific knockout of both AMPK- β subunits,
196 although the mice displayed evidence of dilated cardiomyopathy the rate of glucose and fatty acid
197 oxidation in hearts perfused under normoxic conditions *ex vivo* was normal⁶¹.

198 A final catabolic pathway switched on by AMPK is autophagy, brought about via direct
199 phosphorylation of the protein kinase that triggers that process, ULK1⁶². By triggering digestion of

cellular contents, autophagy may be critical in enhancing cell survival during periods of acute nutrient starvation, such as during ischemia in cardiac muscle. However, it has also been suggested that excessive autophagy might also contribute to cell damage during ischemia/reperfusion⁶³.

Anabolic pathways switched off by AMPK

As well as its classical roles in inhibiting fatty acid and sterol synthesis³⁴, AMPK inactivates key enzymes and regulatory proteins involved in triglyceride/phospholipid synthesis⁶⁴, glycogen synthesis^{13, 14}, rRNA synthesis⁶⁵, and protein synthesis. The latter is inhibited both at the elongation step by phosphorylation and activation of elongation factor-2 (EF2) kinase⁶⁶, and at the initiation step by inactivation of the mechanistic target-of-rapamycin complex-1 (mTORC1) by multiple mechanisms^{67, 68}. By inhibiting mTORC1, AMPK would also be expected to oppose hypertrophy in organs such as the heart, which in most cases is considered to be a deleterious process.

ROLE OF AMPK IN THE HEART

Role in the response to cardiac ischemia

It was reported in 1995 that AMPK was activated by no-flow ischemia in perfused rat hearts, an effect associated with high rates of fatty acid oxidation during reperfusion⁶⁹. AMPK is also activated in the heart by increased workload⁷⁰. The role of AMPK in cardiac ischemia has subsequently been addressed using mouse models where AMPK is down-regulated or absent. The first was a transgenic model where an inactive AMPK- $\alpha 2$ subunit was expressed in both skeletal and cardiac muscle. By competing with endogenous α subunits for available β and γ subunits, the over-expressed inactive $\alpha 2$ subunit down-regulates endogenous $\alpha 1$ and $\alpha 2$, and thus acts as a dominant negative mutant. There was no major phenotype under unstressed conditions, but during low-flow ischemia there was a failure to enhance glucose uptake and glycolysis, while during subsequent reperfusion there was less fatty acid oxidation, lower ATP levels and poorer recovery of contractile function⁷¹. The degree of damage to the myocardium was also greater, suggesting that AMPK exerts a cardioprotective effect overall, even though stimulation of fatty acid oxidation during reperfusion may be deleterious⁶⁹. Using the same model, evidence was obtained that AMPK was required for increased autophagy during ischemia⁷².

227 Next to be studied were conditional knockouts of the upstream kinase LKB1 in both skeletal and
228 cardiac muscle. The basal activity of AMPK- α 2 complexes in the heart was completely abolished,
229 and did not increase in response to no-flow ischemia or anoxia, correlating with a complete lack of
230 ACC phosphorylation at the AMPK site. The AMP:ATP and ADP:ATP ratios were also elevated to
231 a greater extent during no-flow ischemia than in control hearts, confirming that AMPK was
232 protecting the cells against energetic stress. Surprisingly, the activity of AMPK- α 1 complexes was
233 almost unaffected by LKB1 knockout, and still increased in response to ischemia or anoxia⁷³.
234 Broadly similar results were obtained in studies of perfused hearts from mice with a whole body
235 knockout of AMPK- α 2⁷⁴. Interestingly, left ventricular hypertrophy induced by phenylephrine was
236 greatly accentuated in the hearts from AMPK- α 2 knockout mice; this may have been due to less
237 restraint on the mTORC1 pathway, because the basal phosphorylation of p70S6K at the mTORC1
238 site was elevated, although it did not increase further upon phenylephrine treatment as in the
239 controls⁷⁵.

240 ***Mutations in γ 2 and γ 3 subunits causing heart disease and altered glycogen*** 241 ***content***

242 The only mutations in any of the seven genes encoding AMPK subunits clearly shown to cause
243 human pathology are those in *PRKAG2*, encoding AMPK- γ 2. These are associated with multiple
244 cardiac disorders, specifically: (i) ventricular pre-excitation; (ii) excessive glycogen storage in
245 cardiomyocytes; (iii) cardiac hypertrophy. Although rare, the *PRKAG2* mutations are autosomal
246 dominant in effect and therefore occur frequently in affected families. They are invariably missense
247 mutations causing amino acid replacements, of which at least 14 have now been reported (Table 2).
248 The affected residues are perfectly conserved between human γ 1, γ 2 and γ 3, and occur in all four
249 CBS repeats. The location of the residues affected in γ 2, and of the three bound molecules of AMP,
250 are mapped in Fig. 3 onto a structure for γ 1¹⁸. Interestingly, six of the mutations (R302Q, H383R,
251 R384T, H530R, R531G and R531Q) affect basic residues whose side chains directly interact with
252 the phosphate groups of one or more molecules of AMP or ATP, while two others also interact
253 either directly (S548) or indirectly (T400) with AMP¹⁸. Other affected residues lie in more
254 peripheral regions of the γ subunit, and it is less obvious why their replacement should affect
255 function. In the context of bacterially expressed CBS repeats from AMPK- γ 2, the R302Q, H383R,

256 T400N, and R531G mutations all reduced the affinity for ATP as well as AMP, with the severity of
257 the effects increasing in the order R302Q<H383R<T400N<R531G¹⁶. When expressed in
258 mammalian cells as $\alpha 1\beta 1\gamma 2$ heterotrimers, these mutations also reduced allosteric activation by
259 AMP, with the R531G mutation abolishing AMPK activation completely^{16, 76}. The R384T and
260 R531Q mutations were shown later to cause severe effects on AMPK function, similar to or even
261 greater than R531G^{77, 78}. In a comparison of the R531G and R531Q mutations expressed as $\alpha 1\beta 1\gamma 2$
262 heterotrimers in HEK-293 cells, these mutations not only abolished allosteric activation by AMP,
263 but also caused significant increases in basal Thr172 phosphorylation and activity⁷⁸. This was
264 particularly clear in cells stably expressing the R531G mutant, when clones could be selected in
265 which the level of expression of the WT and mutant was identical; the R531G mutant consistently
266 had a 2-fold higher basal Thr172 phosphorylation and activity although, unlike the wild type, it was
267 completely insensitive to further activation by agents that increase cellular AMP/ADP³³.

268 Some of the mutations, such as R302Q, can cause relatively mild symptoms so that the patients
269 may not present in the clinic until early adulthood⁷⁹. By contrast, patients with the R531G mutation
270 develop quite severe symptoms during childhood⁸⁰, while the R384T⁷⁷ and R531Q⁷⁸ mutations were
271 only detected *post mortem* in neonates who had died within weeks of birth, and appeared to be *de*
272 *novo* mutations because neither parent was affected. Consistent with the idea that R531G, R531Q
273 and R384T cause particularly severe forms of the disease, these were the mutations with the largest
274 effects on binding of AMP and ATP to the γ subunit^{16, 77, 78}. It is also interesting that R531 and
275 R384 are involved in the binding of AMP and/or ATP at the critical site, site 3.

276 What causes the cardiac sequelae of the *PRKAG2* mutations? The reduction of AMP binding to
277 the γ subunit causes reduced activation by AMP and is a loss-of-function effect, whereas the
278 reduction of ATP binding may be responsible for the increased basal activity of the R531G and
279 R531Q mutants and is a gain-of-function effect. Since these mutations are dominant, it seems likely
280 that it is the gain-of-function effect, increased basal activity, that causes most of the pathology; the
281 loss-of-function effect may be compensated for by $\gamma 1$, which is the major isoform expressed in
282 heart, at least in rodents⁸¹. This is an important conclusion because pharmaceutical companies are
283 developing AMPK activators to combat Type 2 diabetes, and it suggests that activation of $\gamma 2$ -
284 containing complexes in the heart might have deleterious effects, as seen with these mutations.

285 Why does increased basal activity of γ 2-containing complexes cause these cardiac phenotypes?
286 AMPK is known to promote glucose uptake in cardiac muscle⁷¹, so increased basal activity would
287 be expected to cause a high basal uptake, even in the absence of a real demand for glucose. There is
288 good evidence for this scenario from studies of transgenic mice over-expressing the N488I mutation
289 in the heart, which develop ventricular pre-excitation and cardiac hypertrophy similar to the human
290 disorder⁸². By 50 days of age, they have a 20-fold increase in cardiac glycogen compared with
291 controls, accompanied by higher rates of glucose uptake and glycogen synthesis but lower rates of
292 lactate production, suggesting that increased glucose uptake is being directed into glycogen
293 synthesis rather than glycolysis. Glycogen synthase was also much more highly phosphorylated in
294 the transgenic mice (presumably due to the high basal AMPK activity), but cellular glucose-6-
295 phosphate (G6P, an allosteric activator of glycogen synthase that over-rides the effects of
296 phosphorylation) was also elevated >3-fold. Satisfying confirmation of this interpretation came
297 when the N488I mice were crossed with knock-in mice carrying a mutation that renders glycogen
298 synthase insensitive to G6P. The presence of the G6P-insensitive glycogen synthase reversed the
299 high glycogen phenotype of the N488I mice and rescued the ventricular pre-excitation but not the
300 cardiac hypertrophy, suggesting that the former but not the latter is secondary to increased glycogen
301 content⁸³. Supporting the idea that hyper-activation of the mTORC1 pathway was the cause of
302 hypertrophy, mTORC1 targets such as p70S6K and 4EBP1 were more highly phosphorylated in the
303 N488I mice, while treatment with rapamycin reduced, without completely preventing, the
304 hypertrophy. Increased mTORC1 in the N488I mice is actually rather counter-intuitive, because
305 AMPK is normally thought to inhibit that pathway^{67, 68}. However, a recent study has suggested that
306 AMPK can activate mTORC1 under certain circumstances by sustaining the supply of amino acids
307 via autophagy⁸⁴.

308 Why does abnormally high glycogen content lead to ventricular pre-excitation? During foetal
309 development, the atria and the ventricles become separated by the growth of a fibrous layer called
310 the *annulus fibrosis*, which ensures that the only electrical connection between the two chambers is
311 via the atrioventricular node. In N488I mice the annulus fibrosis appeared to be thin and disrupted
312 in places, and it was suggested that the presence of glycogen-containing vacuoles in myocytes
313 disrupts its formation during fetal development, causing abnormal electrical connections between
314 the atria and ventricles⁸⁵.

Most mouse studies of $\gamma 2$ mutations have involved transgenic mice over-expressing the mutations in human $\gamma 2$. Although these mice do display the key clinical features of the human syndrome, and mice expressing wild type $\gamma 2$ from the same promoters were used as controls, they are not perfect models because $\gamma 2$ is being over-expressed. Recently, three knock-in mouse models, in which mutations in the mouse gene equivalent to human R302Q, N488I or R531G are expressed globally, were studied^{86, 87}. There was evidence for increased basal AMPK activity in liver and/or muscle of all three strains. The N488I and R531G mice displayed ventricular pre-excitation and modest increases in heart weight, which was associated with large increases in glycogen content in hearts of R531G but not the N488I mice. Both N488I and R531G mice displayed resistance to obesity and hepatic steatosis induced by high-fat diet, but, unlike the N488I mice, the R531G mice also displayed impaired renal function associated with glycogen accumulation, cyst formation, and inflammation and apoptosis in the kidney, particularly when on a high-fat diet⁸⁷. By contrast, in the R302Q mice there was no obvious cardiac phenotype, but the homozygotes developed marked obesity as they aged, with smaller effects in heterozygotes. Obesity appeared to be due mainly to increased food intake through enhanced action of ghrelin, whose effects are mediated by AMPK activation via the CaMKK2 pathway in the hypothalamus^{88, 89}. The homozygotes were also hypoinsulemic, apparently due to reduced glucose-stimulated insulin secretion from the pancreas. Overall, the phenotypes of these mice with knock-in mutations in the *PRKAG2* gene showed surprising variability. Interestingly, heterozygous human carriers of the R302Q mutation, who have a relatively mild cardiac phenotype, also display some evidence of increased obesity, as well as higher fasting glucose and glycated haemoglobin (HbA1c) and lower insulin levels than unaffected siblings⁸⁶. These non-cardiac features of the R302Q mutation only became evident following studies of the mouse model.

Finally, all of the residues mutated in AMPK- $\gamma 2$ are also conserved in $\gamma 1$ and $\gamma 3$ (Table 2). It is interesting that none of them have yet been reported to be mutated in the human $\gamma 1$ gene. However, mutations equivalent to R302Q have been found in both pigs and humans in the $\gamma 3$ isoform, which is predominantly expressed in skeletal muscle⁸¹. In human muscle, $\gamma 3$ appears to be present exclusively as the $\alpha 2\beta 2\gamma 3$ complex, which is the only form of AMPK that is activated during exercise⁶. An R200Q mutation (equivalent to R302Q in human $\gamma 2$) was found to be a relatively common dominantly acting genetic variant in Hampshire pigs, and was associated with a high

glycogen content in skeletal muscle; although adversely affecting meat quality, it did not cause obvious clinical problems⁹⁰. Interestingly, in a screen of around 1500 humans, an R225W mutation (R225 being the human equivalent of R302) was found in two apparently unrelated individuals, once again not associated with any obvious clinical defects; muscles from humans with this mutation had a higher basal AMPK activity, 2-fold higher glycogen content and lower triglyceride content⁹¹. In an *in vitro* study of myotubes from R225W carriers, they had 3-fold higher mitochondrial content and oxidative capacity, and 2-fold higher basal glucose uptake and glycogen synthesis rates than matched controls. The R225W subjects also had a remarkable resistance to fatigue during isometric contractions of the quadriceps⁹², suggesting that this mutation might confer an advantage during endurance exercise because of the high glycogen content.

ROLE OF AMPK IN THE VASCULATURE

In addition to the heart, it has become clear that AMPK has an important role in regulating vascular function, with distinct roles in the functions of vascular endothelial cells (ECs), smooth muscle cells (VSMCs), adventitial cells and vascular immune cells.

Regulation of AMPK in endothelial and vascular smooth muscle cells

AMPK- α 1 accounts for the majority of total AMPK activity in ECs⁹³⁻⁹⁵, yet specific down-regulation of AMPK- α 2 still has marked effects^{96, 97}. A wide variety of physiological stimuli have been reported to activate endothelial AMPK, including hypoxia, low glucose and shear stress⁹⁸⁻¹⁰⁰, adiponectin¹⁰¹, angiotensin II and ghrelin^{102, 103}. In addition, CaMKK2 mediates AMPK activation by vasoactive molecules including thrombin, VEGF, sphingosine-1-phosphate, bradykinin and estrogen¹⁰⁴⁻¹⁰⁸, suggesting that any stimulus that increases Ca^{2+} in ECs will activate AMPK. Finally, a number of widely used hypoglycemic (metformin, thiazolidinediones, salicylate, DPP4 inhibitors and liraglutide)¹⁰⁹⁻¹¹² and hypocholesterolemic (statins and fenofibrate)^{113, 114} drugs activate AMPK in cultured ECs.

Physiological signals that *inhibit* AMPK in ECs include high nutrient concentrations¹¹⁵⁻¹¹⁷, yet the mechanisms by which this inhibition of AMPK in ECs occurs are unclear, although increased PP2A-mediated dephosphorylation of AMPK has been proposed¹¹⁵. Endothelial AMPK may therefore be suppressed by the high levels of nutrients associated with obesity and insulin resistance. As protein kinase C (PKC) activation is associated with over-nutrition, it is interesting

that PKC-mediated inhibitory phosphorylation of Ser487 on AMPK- α 1 has been recently reported in ECs¹¹⁸. Conversely, the very first study of AMPK function in ECs demonstrated that AICAR stimulated fatty acid oxidation¹¹⁹ and AMPK activation was subsequently found to normalise impaired fatty acid oxidation and insulin signaling due to high glucose¹²⁰. Stimulation of fat oxidation may partly underlie the effect of AMPK to antagonize palmitate-mediated endothelial dysfunction, thus protecting against lipotoxicity.

As with ECs, AMPK- α 1 accounts for the majority of AMPK activity in murine and human VSMCs^{121, 122}, and low glucose, adiponectin, estradiol and metformin all activate AMPK¹²³⁻¹²⁶. Angiotensin II has also been reported to acutely activate AMPK¹²⁷, although prolonged activation had no effect¹²⁸. Similar to ECs, high glucose inhibits AMPK in VSMCs, and IGF-1 also inhibits AMPK, most likely via inhibitory phosphorylation of AMPK- α 1 by Akt¹²⁹. Several studies have demonstrated that culture in phosphate/ β -glycerophosphate, used to examine VSMC calcification as described later, also inhibits AMPK^{130, 131}.

AMPK and endothelial NO synthesis

Endothelium-derived NO is a key regulator of vascular function, stimulating VSMC relaxation whilst inhibiting pro-inflammatory signaling, leukocyte adhesion, platelet aggregation, VSMC proliferation and migration associated with pathological vascular remodelling and atherosclerosis¹³². The first indication of a specific vascular role for AMPK came when AMPK was shown to phosphorylate at Ser1177 and activate endothelial NO synthase (eNOS), both in cell-free assays and in ECs^{93, 133}. Multiple mechanisms regulate eNOS activity, including Ca²⁺/calmodulin binding, allosteric and protein:protein interactions, co-factor and substrate availability, phosphorylation, and subcellular localisation¹³². In addition to Ser1177, AMPK phosphorylates eNOS at Ser633, required for AICAR-stimulated NO synthesis in HEK293 cells¹³⁴. NO synthesis requires dimerization of eNOS and sufficient tetrahydrobiopterin (BH₄), in whose absence eNOS generates superoxide instead¹³². Another mechanism by which AMPK improves NO synthesis may be by enhancing BH₄ levels, as AMPK prevents degradation of GTP cyclohydrolase I, which is the rate-limiting enzyme in BH₄ synthesis¹³⁵. AMPK can therefore act by multiple mechanisms to stimulate NO production (Fig. 4), although endothelial AMPK activation is not always associated

402 with eNOS phosphorylation or NO synthesis^{94, 136}. AMPK may also mediate some of the
403 endothelial effects of NO, because NO donors activate AMPK in ECs⁹⁶.

404 ***Endothelial AMPK and reactive oxygen species***

405 Inappropriate levels of reactive oxygen species (ROS), in particular superoxide anions generated in
406 response to high concentrations of glucose, lipids and proinflammatory cytokines by NAD(P)H
407 oxidase (Nox), uncoupled eNOS or mitochondrial respiratory chain complexes, have been
408 implicated in vascular disease^{97, 135, 137, 138}. Superoxide reacts with NO to form peroxynitrite,
409 reducing NO bioavailability¹³². AMPK activation in ECs has been widely demonstrated to inhibit
410 ROS formation, increase antioxidant defences and promote mitochondrial biogenesis. Down-
411 regulation of AMPK in ECs increased activity and expression of Nox⁹⁷, whereas AMPK-dependent
412 inhibition of Nox1/2 translocation to the plasma membrane has also been reported¹⁰⁹. The
413 mechanism by which AMPK inhibits Nox remains unclear, but may be secondary to reduced PKC-
414 mediated Nox activation, or NFκB-mediated Nox transcription^{97, 109}. AMPK-dependent inhibition
415 of mitochondrial ROS formation has also been reported in ECs maintained in high glucose¹³⁷,
416 whereas AMPK-mediated stimulation of BH₄ synthesis, as described above, prevents uncoupling of
417 eNOS and superoxide formation¹³⁵. Several groups have reported that AMPK-mediated inhibition
418 of ROS in ECs is associated with increased levels of the antioxidant enzymes superoxide
419 dismutase-2 (SOD2), catalase and thioredoxin^{137, 139, 140}. In ECs, AMPK activation reduces ER
420 stress¹⁴¹, which is tightly linked to oxidative stress and inflammation, whereas silencing of AMPK
421 increases markers of ER stress⁹⁵. Taken together, AMPK activation acts via multiple mechanisms
422 to suppress chronic ROS synthesis in ECs, limiting their damaging actions as well as the
423 sequestration of NO (Fig 4).

424 ***AMPK and vascular cell inflammation***

425 The development of endothelial dysfunction and cardiovascular diseases is associated with elevated
426 TNFα, IL-1β and IL-6¹⁴². TNFα and IL-1β stimulate activation of NFκB and the Jun N-terminal
427 kinase (JNK) pathway, while IL-6 signals via Janus kinases (JAKs) leading to phosphorylation of
428 signal transducer and activation of transcription (STAT) proteins¹⁴³. The anti-inflammatory actions
429 of AMPK in ECs were first described when AMPK activation was shown to suppress palmitate- or
430 TNFα-stimulated NFκB activity¹¹⁷. Under basal conditions, NFκB is in an inactive form in the

cytoplasm due to binding to I κ B whereas, after cytokine stimulation, I κ B phosphorylation by I κ B kinase (IKK) targets it for degradation, allowing NF κ B-mediated transcription of proinflammatory cytokines, adhesion molecules and chemokines^{143, 144}. NO inhibits endothelial NF κ B activity¹⁴⁵, suggesting that AMPK-stimulated NO synthesis would inhibit NF κ B, although NO donors do not effectively suppress NF κ B in ECs with reduced AMPK activity⁹⁶. AMPK- α 2 has been reported to phosphorylate IKK β *in vitro*, inhibiting I κ B phosphorylation and NF κ B activation, with reduced IL-1 β -stimulated IKK phosphorylation observed in ECs lacking AMPK- α 2 but not - α 1⁹⁶. Alternatively, AMPK-mediated phosphorylation of the transcriptional co-activator p300 has been proposed to block acetylation and DNA binding by the p65 subunit of NF κ B¹⁴⁶. The idea that AMPK inhibits NF κ B signalling is reinforced by functional studies demonstrating AMPK-dependent inhibition of NF κ B-regulated expression of adhesion molecules and MCP-1^{96, 144}. Fewer studies have investigated the effect of AMPK on proinflammatory JNK and IL-6 signaling, although AICAR and metformin reduce JNK activity in ECs¹⁴⁷, and increased JNK phosphorylation has been reported in ECs lacking AMPK- α 2⁹⁵. The mechanism of JNK inhibition is uncertain, although AMPK-dependent inhibition of the upstream kinase MKK4 has been reported in other cells¹⁴⁸. Recently, AMPK-dependent inhibition of IL-6-stimulated JAK-STAT signaling has been demonstrated in ECs, potentially via direct inhibitory phosphorylation of JAK1 by AMPK¹¹⁰. AMPK activation therefore appears to rapidly suppress multiple proinflammatory signaling pathways in ECs, by diverse mechanisms.

AMPK activation in VSMCs inhibits TNF α -stimulated NF κ B activity and angiotensin II-stimulated STAT1 activity, as well as reducing expression of inducible NOS and cyclooxygenase-2, and secretion of IL-6 and MCP-1^{149, 150}. Thus, AMPK has anti-inflammatory effects in VSMCs as well as ECs.

AMPK and angiogenesis

Hypoxia, VEGF and adiponectin all stimulate AMPK-dependent EC migration, although there are conflicting reports as to whether this is mediated by NO^{94, 98, 101, 105, 108}. Conversely, down-regulation of AMPK attenuates angiogenesis caused by hypoxia, adiponectin, VEGF or statins, in either tube formation or matrigel plug assays^{94, 98, 101, 113}. Mechanistically, increased UCP2 or SOD2 have been reported to increase angiogenesis in AMPK-deficient ECs^{151, 152}, indicating that down-

460 regulation of ROS may be critical. AMPK activation also stimulates VEGF expression, indicating
461 that AMPK positively influences angiogenesis both by increasing VEGF levels and by increasing
462 VEGF signaling¹⁵³. As angiogenesis would also consume significant amounts of ATP, AMPK
463 activation might also serve a permissive role, ensuring adequate generation of ATP to permit EC
464 migration and proliferation.

465 ***AMPK and VSMC contraction***

466 AMPK has direct anti-contraction effects on VSMCs, because AICAR relaxes aortic rings in an NO-
467 and endothelium-independent manner, an effect lost in AMPK- α 1 knockouts¹²¹. Furthermore,
468 AMPK activation has been reported to affect VSMC contractile signalling, including
469 dephosphorylation of myosin light chain (MLC) and/or myosin phosphatase targeting subunit 1
470 (MYPT1)^{125, 154}. Mechanistically, MLC/MYPT1 dephosphorylation may be a consequence of
471 AMPK-mediated inhibition and phosphorylation of RhoA at Ser188, causing subsequent inhibition
472 of ROCK¹²⁵. A769662 has also been reported to reduce intracellular Ca^{2+} in VSMCs by increasing
473 sarco/endoplasmic Ca^{2+} ATPase (SERCA) activity, associated with increased phosphorylation of
474 phospholamban on Thr17, which disinhibits SERCA and thereby may underlie SERCA activation
475 and vessel relaxation¹⁵⁵.

476 ***AMPK and proliferation, differentiation and migration of VSMCs***

477 Unlike ECs, where AMPK up-regulates proliferation and migration thus supporting angiogenesis, in
478 VSMCs AMPK inhibits proliferation in response to angiotensin II, PDGF and FCS, associated with
479 stimulation of p53 Ser15 phosphorylation and reduced Rb phosphorylation^{128, 156}. VSMCs from
480 mice lacking AMPK- α 2 exhibit increased proliferation, an effect mediated by increased degradation
481 of the cyclin-dependent kinase inhibitor p27^{Kip1} triggered by the ubiquitin E3-ligase Skp2¹⁵⁷.
482 VSMCs retain significant plasticity *in vivo* and can exhibit a synthetic, proliferative phenotype
483 rather than the quiescent, contractile phenotype observed during atherogenesis. AMPK not only
484 suppresses VSMC proliferation, but also inhibits migration and maintains a pro-contraction
485 phenotype^{158, 159}. Inhibition of migration and proliferation are likely to be linked, because increased
486 migration of VSMCs lacking AMPK- α 2 is reported to be Skp2-dependent¹⁵⁹. Furthermore, AICAR
487 limits neointima formation after wire injury of rat femoral arteries, which is likely to reflect the
488 anti-proliferative, anti-migratory actions of AMPK on VSMCs¹²⁷.

489 ***AMPK and VSMC calcification***

490 As mentioned above, culture in high phosphate/ β -glycerophosphate concentrations is used
491 experimentally to stimulate calcium deposition in VSMCs. *In vivo*, such vascular calcification is
492 frequently associated with ageing, atherosclerosis and diabetes mellitus. AICAR, adiponectin and
493 metformin all inhibit VSMC calcification *in vitro*^{130, 131, 160}. In recent studies of atherosclerosis-
494 prone *ApoE*^{-/-} mice, deletion of AMPK- α 1 but not AMPK- α 2 caused greater calcification of
495 atherogenic plaques, and levels of the osteogenic transcription factor Runx2¹⁶⁰. Furthermore,
496 metformin reduced atherosclerotic calcification and Runx2 expression in the mice, an effect that
497 was absent in *ApoE*^{-/-} mice lacking AMPK- α 1. The authors of that study further demonstrated that
498 VSMC-specific AMPK- α 1 deletion in *ApoE*^{-/-} mice phenocopied the increased calcification and
499 Runx2 expression, whereas macrophage-specific AMPK- α 1 deletion had no effect. The mechanism
500 underlying the AMPK-mediated inhibition of Runx2 levels was further proposed to be mediated by
501 phosphorylation of PIAS1 (protein inhibitor of activated STAT-1) at Ser510, which acts as a
502 SUMO E3-ligase to trigger Runx2 SUMOylation and degradation¹⁶⁰.

503 **ROLE OF AMPK IN THE VASCULATURE IN VIVO**

504 As many of the actions of AMPK in ECs and VSMCs should have beneficial effects on vascular
505 function (Fig 5), considerable efforts have been made to examine whether AMPK influences
506 vascular tone, remodelling and atherogenesis *in vivo*. Under physiological conditions, vascular
507 AMPK has been reported to be activated *in vivo* by exercise¹⁶¹ and estradiol¹²⁵, and is suppressed by
508 high fat and fructose diets in rodents^{162, 163}. Furthermore, exercise stimulates aortic AMPK
509 phosphorylation in both ECs and VSMCs as assessed by immunohistochemistry¹⁶¹.

510 ***AMPK and vascular tone***

511 As described above, AMPK can stimulate NO synthesis by ECs and independently inhibit
512 contractile protein function in VSMCs *in vitro*. In intact arterial vessels, AICAR stimulates
513 vasodilation in a diverse range of vascular beds from several species^{121, 164-167}, an effect greatly
514 attenuated in mice lacking AMPK- α 1¹²¹. The mechanism of AICAR-stimulated vasodilation
515 remains uncertain, and has been variously reported to be endothelium- and NO-dependent^{164, 167},
516 partially NO- and endothelium-dependent^{165, 166} or NO- and endothelium-independent¹²¹. Using
517 endothelium-specific knockouts it has been proposed that AMPK- α 1 is important for endothelium-

518 dependent hyperpolarisation-mediated relaxation of resistance arteries¹⁶⁸. Interestingly, resistance
519 arteries exhibited endothelium-independent dilation in response to A769662¹⁵⁵, suggesting a
520 VSMC-mediated effect. It is possible that ECs and VSMCs exhibit differential sensitivities to
521 AICAR and A769662, or that some of these may be AMPK-independent effects. Despite studies
522 indicating that AMPK- α 2 has a less important role in regulation of vascular tone^{121, 168}, AMPK- α 2
523 knockout mice are hypertensive and exhibit increased contractile responses to phenylephrine in
524 aortic rings¹⁶⁹. Furthermore, impaired bradykinin-dependent vasodilation has been described in EC-
525 specific AMPK- α 2 knockout mice, thought to be due to increased bradykinin degradation by
526 angiotensin-converting enzyme activity¹⁷⁰. AICAR rapidly reduced blood pressure in spontaneously
527 hypertensive rats, but was without effect in normotensive controls¹⁶⁵, while prolonged
528 administration of AICAR reduced systolic blood pressure in obese Zucker rats¹⁷¹. These data
529 support a role for AMPK in the regulation of vascular tone in disease models, but cannot exclude
530 systemic actions on the heart, kidney or other tissues. Surprisingly, whether systemic administration
531 of more selective AMPK activators, such as A769662, 991 or C13, alters vascular tone has yet to be
532 reported. More recently it has become clear that perivascular adipose tissue (PVAT), which is
533 removed in most myography protocols, has a paracrine anti-contractile effect on the underlying
534 vessel. AMPK activity in resistance arteries has been shown to alter the influence of PVAT-derived
535 mediators¹⁷² and the anti-contractile action of PVAT is absent in mice lacking AMPK- α 1, perhaps
536 due to reduced adiponectin secretion¹⁷³. AMPK in ECs, VSMCs and adventitial PVAT may
537 therefore all contribute to the maintenance of vascular tone, but the cell type that mediates actions
538 of AMPK on vascular tone may change, depending on the location of the vessel.

539 ***AMPK and atherosclerosis***

540 The potential anti-atherogenic actions of AMPK have been investigated in vascular injury models
541 and atherosclerosis-prone hypercholesterolemic mice. AICAR attenuates post-ischaemic leukocyte
542 rolling and adhesion to the endothelium *in vivo*, an effect lost in AMPK- α 1 or AMPK- α 2 knockout
543 mice¹⁷⁴, and systemic administration of AICAR or metformin reduces atherosclerotic lesion size,
544 macrophage accumulation and inflammation¹⁷⁵⁻¹⁷⁷. Similarly, berberine reduced the severity of
545 atherosclerotic lesions in atherosclerosis-prone mice, an effect attenuated in AMPK- α 2
546 knockouts¹⁷⁸. AMPK α 2-deficient mice also exhibited increased atherosclerosis⁹⁵, although the

547 global deficiency makes it difficult to assess whether this was due to a direct effect on vascular
548 tissues. With respect to plaque stability and progression, recent studies have yielded exciting
549 results. Metformin reduced plaque calcification, and mice with a SMC-specific knockout of
550 AMPK- α 1 had exacerbated calcification of atherosclerotic plaques in brachiocephalic arteries,
551 phenocopying the increased calcification observed in global AMPK- α 1 but not AMPK- α 2
552 knockouts¹⁶⁰. As the clinical use of metformin is associated with reduced macrovascular morbidity
553 and mortality independently of glycemia¹⁷⁹, these results may help to define the pathways involved.
554 Using a ligation model for studying injury-induced neointima stability in brachiocephalic arteries,
555 mice with deletion of AMPK- α 1 but not AMPK- α 2 exhibited more occlusive lesions, with lower
556 collagen and higher macrophage content, indicative of plaque instability¹⁸⁰. Similarly, in a high fat
557 diet model, SMC-specific AMPK- α 2 knockouts exhibited features of unstable plaques, including
558 phenotype switching of VSMCs¹⁵⁸. These data therefore indicate that AMPK activation may not
559 only inhibit atherogenesis, but also inhibit the generation of vulnerable, calcified plaques. Repair of
560 damaged endothelium is also considered important to prevent endothelial dysfunction after injury,
561 and EC-specific expression of constitutively active AMPK has also been reported to promote re-
562 endothelialization in a wire injury model, attributable in part to increased mobilization and
563 incorporation of endothelial progenitor cells¹⁸¹.

564 Immune cells are involved in all stages of atherosclerosis, and AMPK suppresses inflammatory
565 signalling, monocyte-to-macrophage differentiation and foam cell formation^{175, 177, 182}. Indeed,
566 myeloid-specific AMPK- α 1 knockout mice fed an atherogenic diet on a LDL receptor knockout
567 background have recently been shown to exhibit exacerbated atherosclerosis, with increased plaque
568 macrophage content and inflammatory gene expression¹⁸². All of these studies suggest that AMPK
569 activation limits atherosclerosis, although all of the results are from rodent models, rather than
570 humans. One intriguing question is how, despite only contributing a small fraction of total AMPK
571 activity in vascular cells, specific down-regulation of AMPK- α 2 has such marked effects on
572 atheroma development in mice^{95, 158, 174, 178}, similar to the specific effects noted with respect to
573 vascular tone^{169, 170}. It remains to be determined whether there are isoform-specific substrates or
574 specific subcellular localisations that contribute to the vascular function of AMPK- α 2.

575 ***AMPK and pulmonary vascular remodelling***

576 Several studies have identified a critical role for AMPK in the pulmonary vascular remodelling and
577 perivascular inflammation that characterizes pulmonary arterial hypertension (PAH). Metformin
578 reduces pulmonary artery VSMC proliferation in an AMPK-dependent manner¹⁸³, and inhibits the
579 vascular remodelling of pulmonary arteries in rodent models of PAH^{184, 185}. Mice with an EC-
580 specific lack of AMPK also exhibit accelerated pulmonary remodeling¹⁸⁵. Whether the beneficial
581 action of metformin in PAH is mediated by AMPK remains to be determined. By contrast, AMPK
582 activation by hypoxia may contribute to the development of PAH by promoting pulmonary VSMC
583 survival¹⁸⁶, while AMPK-mediated inhibition of the voltage-gated K⁺ channel Kv1.5 may underlie
584 the acute detrimental effects of hypoxia on PAH¹⁸⁷.

585 ***Conclusions and Perspectives***

586 AMPK exists as heterotrimeric complexes consisting of catalytic α subunits and regulatory β and γ
587 subunits. AMPK complexes sense the energy status of cells by sensing increases in the cellular
588 AMP:ATP and/or ADP:ATP ratios. AMP, ADP and ATP bind at up to three sites on the γ subunits;
589 binding of AMP and/or ADP causes activation by promoting net phosphorylation at Thr172 within
590 the activation loop on the α subunit, while binding of AMP only causes further allosteric activation.
591 Once activated by energy stress, AMPK switches on catabolic pathways that generate ATP, while
592 switching off cell growth and proliferation and other processes that consume ATP.

593 AMPK appears to exert a protective effect in rodent heart during ischemic episodes. Inherited
594 and/or *de novo* mutations in the *PRKAG2* gene (encoding AMPK- γ 2) in humans cause heart disease
595 of varying severity characterized by ventricular pre-excitation, excessive cardiac glycogen content,
596 and hypertrophy. The mutations cause an increase in basal AMPK activity, leading to increased
597 glucose uptake that accounts for the first two abnormalities. They also cause a failure of γ 2-
598 containing complexes to be further activated by AMP, which might explain the hypertrophy
599 through unrestrained activity of the mTORC1 pathway.

600 In blood vessels, AMPK inactivation is associated with anti-contractile, anti-inflammatory and
601 anti-atherogenic actions on both the vascular endothelium and smooth muscle. Exciting recent
602 studies link AMPK activation to increased stability and reduced calcification of atherosclerotic
603 plaques as well as highlighting a potential role for AMPK in pulmonary arterial hypertension.

604 Given the critical role of AMPK in the regulation of nutrient metabolism, therapies that activate
605 AMPK may not only normalise metabolic dysfunction but also reduce the burden of cardiovascular
606 complications in obesity and type 2 diabetes. Recent studies using vascular tissue-specific deletion
607 of AMPK- α isoforms are beginning to elucidate specific roles for AMPK- α isoforms, yet despite
608 the wealth of research demonstrating the functional cardiovascular consequences of AMPK
609 activation, the AMPK substrates involved and underlying mechanisms in several cases remain
610 poorly defined. Emerging technologies such as phosphoproteomics may prove beneficial in
611 understanding such mechanisms after AMPK up- or down-regulation in cardiovascular tissues in
612 different disease settings and disease models.

613 ***Acknowledgements***

614 Recent studies in the DGH laboratory have been supported by a Senior Investigator Award from the
615 Wellcome Trust (097726) and a Programme Grant from Cancer Research UK (C37030/A15101).
616 Recent studies in the IPS laboratory have been supported by Project Grants from the British Heart
617 Foundation (PG/12/1/29276 and PG/13/82/30483) and Diabetes UK (BDA09/0003904 and
618 BDA12/0004489).

619 ***References***

- 620 1. Hardie DG, Schaffer BE, Brunet A. AMPK: an energy-sensing pathway with multiple
621 inputs and outputs. *Trends Cell Biol.* 2016;26:190-201
- 622 2. Hardie DG, Ashford ML. AMPK: regulating energy balance at the cellular and whole body
623 levels. *Physiology.* 2014;29:99-107
- 624 3. Hardie DG, Ross FA, Hawley SA. AMPK: a nutrient and energy sensor that maintains
625 energy homeostasis. *Nature Rev. Mol. Cell Biol.* 2012;13:251-262
- 626 4. Daskalopoulos EP, Dufeys C, Beauloye C, Bertrand L, Horman S. AMPK in cardiovascular
627 diseases. *EXS.* 2016;107:179-201
- 628 5. Daskalopoulos EP, Dufeys C, Bertrand L, Beauloye C, Horman S. AMPK in cardiac fibrosis
629 and repair: Actions beyond metabolic regulation. *J. Mol. Cell. Cardiol.* 2016;91:188-200
- 630 6. Birk JB, Wojtaszewski JF. Predominant α -1/ β -2/ γ -3 AMPK activation
631 during exercise in human skeletal muscle. *J. Physiol.* 2006;577:1021-1032
- 632 7. Li X, Wang L, Zhou XE, Ke J, de Waal PW, Gu X, Tan MH, Wang D, Wu D, Xu HE, Melcher K.
633 Structural basis of AMPK regulation by adenine nucleotides and glycogen. *Cell Res.*
634 2015;25:50-66

- 635 8. Xiao B, Sanders MJ, Carmena D, Bright NJ, Haire LF, Underwood E, Patel BR, Heath RB,
636 Walker PA, Hallen S, Giordanetto F, Martin SR, Carling D, Gamblin SJ. Structural basis of
637 AMPK regulation by small molecule activators. *Nature Commun.* 2013;4:3017
- 638 9. Calabrese MF, Rajamohan F, Harris MS, Caspers NL, Magyar R, Withka JM, Wang H,
639 Borzilleri KA, Sahasrabudhe PV, Hoth LR, Geoghegan KF, Han S, Brown J, Subashi TA,
640 Reyes AR, Frisbie RK, Ward J, Miller RA, Landro JA, Londregan AT, Carpino PA, Cabral S,
641 Smith AC, Conn EL, Cameron KO, Qiu X, Kurumbail RG. Structural basis for AMPK
642 activation: natural and synthetic ligands regulate kinase activity from opposite poles by
643 different molecular mechanisms. *Structure.* 2014;22:1161-1172
- 644 10. Hudson ER, Pan DA, James J, Lucocq JM, Hawley SA, Green KA, Baba O, Terashima T,
645 Hardie DG. A novel domain in AMP-activated protein kinase causes glycogen storage
646 bodies similar to those seen in hereditary cardiac arrhythmias. *Curr. Biol.* 2003;13:861-
647 866
- 648 11. Polekhina G, Gupta A, Michell BJ, van Denderen B, Murthy S, Feil SC, Jennings IG,
649 Campbell, D.J., Witters LA, Parker MW, Kemp BE, Stapleton D. AMPK b-Subunit targets
650 metabolic stress-sensing to glycogen. *Current Biol.* 2003;13:867-871
- 651 12. Bendayan M, Londono I, Kemp BE, Hardie GD, Ruderman N, Prentki M. Association of
652 AMP-activated protein kinase subunits with glycogen particles as revealed in situ by
653 immunoelectron microscopy. *J. Histochem. Cytochem.* 2009;57:963-971
- 654 13. Jorgensen SB, Nielsen JN, Birk JB, Olsen GS, Viollet B, Andreelli F, Schjerling P, Vaulont S,
655 Hardie DG, Hansen BF, Richter EA, Wojtaszewski JF. The α 2-5'AMP-activated protein
656 kinase is a site 2 glycogen synthase kinase in skeletal muscle and is responsive to
657 glucose loading. *Diabetes.* 2004;53:3074-3081
- 658 14. Bultot L, Guigas B, Von Wilamowitz-Moellendorff A, Maisin L, Vertommen D, Hussain N,
659 Beullens M, Guinovart JJ, Foretz M, Viollet B, Sakamoto K, Hue L, Rider MH. AMP-
660 activated protein kinase phosphorylates and inactivates liver glycogen synthase.
661 *Biochem. J.* 2012;443:193-203
- 662 15. Langendorf CG, Kemp BE. Choreography of AMPK activation. *Cell Res.* 2015;25:5-6
- 663 16. Scott JW, Hawley SA, Green KA, Anis M, Stewart G, Scullion GA, Norman DG, Hardie DG.
664 CBS domains form energy-sensing modules whose binding of adenosine ligands is
665 disrupted by disease mutations. *J. Clin. Invest.* 2004;113:274-284
- 666 17. Townley R, Shapiro L. Crystal structures of the adenylate sensor from fission yeast AMP-
667 activated protein kinase. *Science.* 2007;315:1726-1729
- 668 18. Xiao B, Heath R, Saiu P, Leiper FC, Leone P, Jing C, Walker PA, Haire L, Eccleston JF, Davis
669 CT, Martin SR, Carling D, Gamblin SJ. Structural basis for AMP binding to mammalian
670 AMP-activated protein kinase. *Nature.* 2007;449:496-500
- 671 19. Oakhill JS, Chen ZP, Scott JW, Steel R, Castelli LA, Ling N, Macaulay SL, Kemp BE. beta-
672 Subunit myristoylation is the gatekeeper for initiating metabolic stress sensing by AMP-
673 activated protein kinase (AMPK). *Proc. Natl. Acad. Sci. USA.* 2010;107:19237-19241

- 674 20. Hong SP, Leiper FC, Woods A, Carling D, Carlson M. Activation of yeast Snf1 and
675 mammalian AMP-activated protein kinase by upstream kinases. *Proc. Natl. Acad. Sci. U S*
676 *A*. 2003;100:8839-8843
- 677 21. Sutherland CM, Hawley SA, McCartney RR, Leech A, Stark MJ, Schmidt MC, Hardie DG.
678 Elm1p is one of three upstream kinases for the *Saccharomyces cerevisiae* SNF1 complex.
679 *Curr. Biol*. 2003;13:1299-1305
- 680 22. Woods A, Johnstone SR, Dickerson K, Leiper FC, Fryer LG, Neumann D, Schlattner U,
681 Wallimann T, Carlson M, Carling D. LKB1 is the upstream kinase in the AMP-activated
682 protein kinase cascade. *Curr. Biol*. 2003;13:2004-2008
- 683 23. Hawley SA, Boudeau J, Reid JL, Mustard KJ, Udd L, Makela TP, Alessi DR, Hardie DG.
684 Complexes between the LKB1 tumor suppressor, STRADa/b and MO25a/b are upstream
685 kinases in the AMP-activated protein kinase cascade. *J. Biol*. 2003;2:28
- 686 24. Shaw RJ, Kosmatka M, Bardeesy N, Hurley RL, Witters LA, DePinho RA, Cantley LC. The
687 tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates
688 apoptosis in response to energy stress. *Proc. Natl. Acad. Sci. USA*. 2004;101:3329-3335
- 689 25. Hawley SA, Pan DA, Mustard KJ, Ross L, Bain J, Edelman AM, Frenguelli BG, Hardie DG.
690 Calmodulin-dependent protein kinase kinase-beta is an alternative upstream kinase for
691 AMP-activated protein kinase. *Cell Metab*. 2005;2:9-19
- 692 26. Woods A, Dickerson K, Heath R, Hong SP, Momcilovic M, Johnstone SR, Carlson M,
693 Carling D. Ca²⁺/calmodulin-dependent protein kinase kinase-beta acts upstream of
694 AMP-activated protein kinase in mammalian cells. *Cell Metab*. 2005;2:21-33
- 695 27. Hurley RL, Anderson KA, Franzone JM, Kemp BE, Means AR, Witters LA. The
696 Ca²⁺/calmodulin-dependent protein kinase kinases are AMP-activated protein kinase
697 kinases. *J. Biol. Chem*. 2005;280:29060-29066
- 698 28. Alessi DR, Sakamoto K, Bayascas JR. Lkb1-dependent signaling pathways. *Annu. Rev.*
699 *Biochem*. 2006;75:137-163
- 700 29. Hardie DG, Hawley SA. AMP-activated protein kinase: the energy charge hypothesis
701 revisited. *BioEssays*. 2001;23:1112-1119
- 702 30. Gowans GJ, Hawley SA, Ross FA, Hardie DG. AMP is a true physiological regulator of
703 AMP-activated protein kinase by both allosteric activation and enhancing net
704 phosphorylation. *Cell Metab*. 2013;18:556-566
- 705 31. Ross FA, Jensen TE, Hardie DG. Differential regulation by AMP and ADP of AMPK
706 complexes containing different gamma subunit isoforms. *Biochem. J*. 2016;473:189-199
- 707 32. Xiao B, Sanders MJ, Underwood E, Heath R, Mayer FV, Carmena D, Jing C, Walker PA,
708 Eccleston JF, Haire LF, Saiu P, Howell SA, Aasland R, Martin SR, Carling D, Gamblin SJ.
709 Structure of mammalian AMPK and its regulation by ADP. *Nature*. 2011;472:230-233
- 710 33. Hawley SA, Ross FA, Chevtzoff C, Green KA, Evans A, Fogarty S, Towler MC, Brown LJ,
711 Ogunbayo OA, Evans AM, Hardie DG. Use of cells expressing gamma subunit variants to
712 identify diverse mechanisms of AMPK activation. *Cell Metab*. 2010;11:554-565

- 713 34. Corton JM, Gillespie JG, Hawley SA, Hardie DG. 5-Aminoimidazole-4-carboxamide
714 ribonucleoside: a specific method for activating AMP-activated protein kinase in intact
715 cells? *Eur. J. Biochem.* 1995;229:558-565
- 716 35. Longnus SL, Wambolt RB, Parsons HL, Brownsey RW, Allard MF. 5-Aminoimidazole-4-
717 carboxamide 1-beta -D-ribofuranoside (AICAR) stimulates myocardial glycogenolysis by
718 allosteric mechanisms. *Am. J. Physiol.* 2003;284:R936-R944
- 719 36. Vincent MF, Marangos PJ, Gruber HE, Van den Berghe G. Inhibition by AICA riboside of
720 gluconeogenesis in isolated rat hepatocytes. *Diabetes.* 1991;40:1259-1266
- 721 37. Gadalla AE, Pearson T, Currie AJ, Dale N, Hawley SA, Sheehan M, Hirst W, Michel AD,
722 Randall A, Hardie DG, Frenguelli BG. AICA riboside both activates AMP-activated protein
723 kinase and competes with adenosine for the nucleoside transporter in the CA1 region of
724 the rat hippocampus. *J. Neurochem.* 2004;88:1272-1282
- 725 38. Gomez-Galeno JE, Dang Q, Nguyen TH, Boyer SH, Grote MP, Sun Z, Chen M, Craigo WA,
726 van Poelje PD, MacKenna DA, Cable EE, Rolzin PA, Finn PD, Chi B, Linemeyer DL, Hecker
727 SJ, Erion MD. A potent and selective AMPK activator that inhibits de novo lipogenesis.
728 *ACS Med. Chem. Lett.* 2010;1:478-482
- 729 39. Hunter RW, Foretz M, Bultot L, Fullerton MD, Deak M, Ross FA, Hawley SA, Shpiro N,
730 Viollet B, Barron D, Kemp BE, Steinberg GR, Hardie DG, Sakamoto K. Mechanism of
731 action of Compound-13: an alpha1-selective small molecule activator of AMPK. *Chem.*
732 *Biol.* 2014;21:866-879
- 733 40. Langendorf CG, Ngoei KR, Scott JW, Ling NX, Issa SM, Gorman MA, Parker MW, Sakamoto
734 K, Oakhill JS, Kemp BE. Structural basis of allosteric and synergistic activation of AMPK
735 by furan-2-phosphonic derivative C2 binding. *Nat. Commun.* 2016;7:10912
- 736 41. Sanders MJ, Ali ZS, Hegarty BD, Heath R, Snowden MA, Carling D. Defining the
737 mechanism of activation of AMP-activated protein kinase by the small molecule A-
738 769662, a member of the thienopyridone family. *J. Biol. Chem.* 2007;282:32539-32548
- 739 42. Goransson O, McBride A, Hawley SA, Ross FA, Shpiro N, Foretz M, Viollet B, Hardie DG,
740 Sakamoto K. Mechanism of action of A-769662, a valuable tool for activation of AMP-
741 activated protein kinase. *J. Biol. Chem.* 2007;282:32549-32560
- 742 43. Scott JW, Ling N, Issa SM, Dite TA, O'Brien MT, Chen ZP, Galic S, Langendorf CG, Steinberg
743 GR, Kemp BE, Oakhill JS. Small molecule drug A-769662 and AMP synergistically activate
744 naive AMPK independent of upstream kinase signaling. *Chem. Biol.* 2014;21:619-627
- 745 44. Hawley SA, Fullerton MD, Ross FA, Schertzer JD, Chevtzoff C, Walker KJ, Pegg MW,
746 Zibrova D, Green KA, Mustard KJ, Kemp BE, Sakamoto K, Steinberg GR, Hardie DG. The
747 ancient drug salicylate directly activates AMP-activated protein kinase. *Science.*
748 2012;336:918-922
- 749 45. Bain J, Plater L, Elliott M, Shpiro N, Hastie CJ, McLauchlan H, Klevernic I, Arthur JS, Alessi
750 DR, Cohen P. The selectivity of protein kinase inhibitors: a further update. *Biochem J.*
751 2007;408:297-315

- 752 46. Wu N, Zheng B, Shaywitz A, Dagon Y, Tower C, Bellinger G, Shen CH, Wen J, Asara J,
753 McGraw TE, Kahn BB, Cantley LC. AMPK-dependent degradation of TXNIP upon energy
754 stress leads to enhanced glucose uptake via GLUT1. *Mol. Cell.* 2013;49:1167-1175
- 755 47. Chen Q, Xie B, Zhu S, Rong P, Sheng Y, Ducommun S, Chen L, Quan C, Li M, Sakamoto K,
756 MacKintosh C, Chen S, Wang HY. A Tbc1d1 Ser231Ala-knockin mutation partially
757 impairs AICAR- but not exercise-induced muscle glucose uptake in mice. *Diabetologia.*
758 2017;60:336-345
- 759 48. Zheng D, MacLean PS, Pohnert SC, Knight JB, Olson AL, Winder WW, Dohm GL.
760 Regulation of muscle GLUT-4 transcription by AMP-activated protein kinase. *J. Appl.*
761 *Physiol.* 2001;91:1073-1083
- 762 49. McGee SL, van Denderen BJ, Howlett KF, Mollica J, Schertzer JD, Kemp BE, Hargreaves M.
763 AMP-activated protein kinase regulates GLUT4 transcription by phosphorylating histone
764 deacetylase 5. *Diabetes.* 2008;57:860-867
- 765 50. Mihaylova MM, Vasquez DS, Ravnskjaer K, Denechaud PD, Yu RT, Alvarez JG, Downes M,
766 Evans RM, Montminy M, Shaw RJ. Class IIa histone deacetylases are hormone-activated
767 regulators of FOXO and mammalian glucose homeostasis. *Cell.* 2011;145:607-621
- 768 51. Marsin AS, Bertrand L, Rider MH, Deprez J, Beauloye C, Vincent MF, Van den Berghe G,
769 Carling D, Hue L. Phosphorylation and activation of heart PFK-2 by AMPK has a role in
770 the stimulation of glycolysis during ischaemia. *Current Biol.* 2000;10:1247-1255
- 771 52. Marsin AS, Bouzin C, Bertrand L, Hue L. The stimulation of glycolysis by hypoxia in
772 activated monocytes is mediated by AMP-activated protein kinase and inducible 6-
773 phosphofructo-2-kinase. *J. Biol. Chem.* 2002;277:30778-30783
- 774 53. Rider MH, Bertrand L, Vertommen D, Michels PA, Rousseau GG, Hue L. 6-phosphofructo-
775 2-kinase/fructose-2,6-bisphosphatase: head-to-head with a bifunctional enzyme that
776 controls glycolysis. *Biochem. J.* 2004;381:561-579
- 777 54. O'Neill HM, Lally JS, Galic S, Thomas M, Azizi PD, Fullerton MD, Smith BK, Puliniikunnil T,
778 Chen Z, Samaan MC, Jorgensen SB, Dyck JR, Holloway GP, Hawke TJ, van Denderen BJ,
779 Kemp BE, Steinberg GR. AMPK phosphorylation of ACC2 is required for skeletal muscle
780 fatty acid oxidation and insulin sensitivity in mice. *Diabetologia.* 2014;57:1693-1702
- 781 55. Merrill GM, Kurth E, Hardie DG, Winder WW. AICAR decreases malonyl-CoA and
782 increases fatty acid oxidation in skeletal muscle of the rat. *Am. J. Physiol.*
783 1997;273:E1107-E1112
- 784 56. Fullerton MD, Galic S, Marcinko K, Sikkema S, Puliniikunnil T, Chen ZP, O'Neill HM, Ford
785 RJ, Palanivel R, O'Brien M, Hardie DG, Macaulay SL, Schertzer JD, Dyck JR, van Denderen
786 BJ, Kemp BE, Steinberg GR. Single phosphorylation sites in ACC1 and ACC2 regulate lipid
787 homeostasis and the insulin-sensitizing effects of metformin. *Nat. Med.* 2013;19:1649-
788 1654
- 789 57. Winder WW, Holmes BF, Rubink DS, Jensen EB, Chen M, Holloszy JO. Activation of AMP-
790 activated protein kinase increases mitochondrial enzymes in skeletal muscle. *J. Appl.*
791 *Physiol.* 2000;88:2219-2226

- 792 58. Jager S, Handschin C, St-Pierre J, Spiegelman BM. AMP-activated protein kinase (AMPK)
793 action in skeletal muscle via direct phosphorylation of PGC-1{alpha}. *Proc. Natl. Acad.*
794 *Sci. USA.* 2007;104:12017-12022
- 795 59. Canto C, Gerhart-Hines Z, Feige JN, Lagouge M, Noriega L, Milne JC, Elliott PJ, Puigserver
796 P, Auwerx J. AMPK regulates energy expenditure by modulating NAD(+) metabolism and
797 SIRT1 activity. *Nature.* 2009;458:1056-1060
- 798 60. Lin J, Handschin C, Spiegelman BM. Metabolic control through the PGC-1 family of
799 transcription coactivators. *Cell Metab.* 2005;1:361-370
- 800 61. Sung MM, Zordoky BN, Bujak AL, Lally JS, Fung D, Young ME, Horman S, Miller EJ, Light
801 PE, Kemp BE, Steinberg GR, Dyck JR. AMPK deficiency in cardiac muscle results in
802 dilated cardiomyopathy in the absence of changes in energy metabolism. *Cardiovasc. Res.*
803 2015;107:235-245
- 804 62. Egan DF, Shackelford DB, Mihaylova MM, Gelino S, Kohnz RA, Mair W, Vasquez DS, Joshi
805 A, Gwinn DM, Taylor R, Asara JM, Fitzpatrick J, Dillin A, Viollet B, Kundu M, Hansen M,
806 Shaw RJ. Phosphorylation of ULK1 (hATG1) by AMP-activated protein kinase connects
807 energy sensing to mitophagy. *Science.* 2011;331:456-461
- 808 63. Wang S, Wang C, Yan F, Wang T, He Y, Li H, Xia Z, Zhang Z. N-Acetylcysteine attenuates
809 diabetic myocardial ischemia reperfusion injury through inhibiting excessive autophagy.
810 *Mediators Inflamm.* 2017;2017:9257291
- 811 64. Muoio DM, Seefeld K, Witters LA, Coleman RA. AMP-activated kinase reciprocally
812 regulates triacylglycerol synthesis and fatty acid oxidation in liver and muscle: evidence
813 that sn-glycerol- 3-phosphate acyltransferase is a novel target. *Biochem. J.*
814 1999;338:783-791
- 815 65. Hoppe S, Bierhoff H, Cado I, Weber A, Tiebe M, Grummt I, Voit R. AMP-activated protein
816 kinase adapts rRNA synthesis to cellular energy supply. *Proc. Natl. Acad. Sci. USA.*
817 2009;106:17781-17786
- 818 66. Browne GJ, Finn SG, Proud CG. Stimulation of the AMP-activated protein kinase leads to
819 activation of eukaryotic elongation factor 2 kinase and to its phosphorylation at a novel
820 site, serine 398. *J. Biol. Chem.* 2004;279:12220-12231
- 821 67. Inoki K, Zhu T, Guan KL. TSC2 mediates cellular energy response to control cell growth
822 and survival. *Cell.* 2003;115:577-590
- 823 68. Gwinn DM, Shackelford DB, Egan DF, Mihaylova MM, Mery A, Vasquez DS, Turk BE, Shaw
824 RJ. AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol. Cell.*
825 2008;30:214-226
- 826 69. Kudo N, Barr AJ, Barr RL, Desai S, Lopaschuk GD. High rates of fatty acid oxidation
827 during reperfusion of ischemic hearts are associated with a decrease in malonyl-CoA
828 levels due to an increase in 5'-AMP-activated protein kinase inhibition of acetyl-CoA
829 carboxylase. *J. Biol. Chem.* 1995;270:17513-17520

- 830 70. Coven DL, Hu X, Cong L, Bergeron R, Shulman GI, Hardie DG, Young LH. Physiologic role
831 of AMP-activated protein kinase (AMPK) in the heart: graded activation during exercise.
832 *Am. J. Physiol.* 2003;285:E629-E636
- 833 71. Russell RR, 3rd, Li J, Coven DL, Pypaert M, Zechner C, Palmeri M, Giordano FJ, Mu J,
834 Birnbaum MJ, Young LH. AMP-activated protein kinase mediates ischemic glucose
835 uptake and prevents postischemic cardiac dysfunction, apoptosis, and injury. *J. Clin.*
836 *Invest.* 2004;114:495-503
- 837 72. Matsui Y, Takagi H, Qu X, Abdellatif M, Sakoda H, Asano T, Levine B, Sadoshima J. Distinct
838 roles of autophagy in the heart during ischemia and reperfusion: roles of AMP-activated
839 protein kinase and Beclin 1 in mediating autophagy. *Circ. Res.* 2007;100:914-922
- 840 73. Sakamoto K, Zarrinpashneh E, Budas GR, Pouleur AC, Dutta A, Prescott AR,
841 Vanoverschelde JL, Ashworth A, Jovanovic A, Alessi DR, Bertrand L. Deficiency of LKB1
842 in heart prevents ischemia-mediated activation of AMPK α 2 but not AMPK α 1.
843 *Am. J. Physiol. Endocrinol. Metab.* 2006;290:E780-E788
- 844 74. Zarrinpashneh E, Carjaval K, Beauloye C, Ginion A, Mateo P, Pouleur AC, Horman S,
845 Vaulont S, Hoerter J, Viollet B, Hue L, Vanoverschelde JL, Bertrand L. Role of the α 2-
846 isoform of AMP-activated protein kinase in the metabolic response of the heart to no-
847 flow ischemia. *Am. J. Physiol. Heart Circ. Physiol.* 2006;291:H2875-H2883
- 848 75. Zarrinpashneh E, Beauloye C, Ginion A, Pouleur AC, Havaux X, Hue L, Viollet B,
849 Vanoverschelde JL, Bertrand L. AMPK- α 2 counteracts the development of cardiac
850 hypertrophy induced by isoproterenol. *Biochem. Biophys. Res. Commun.* 2008;376:677-
851 681
- 852 76. Daniel T, Carling D. Functional analysis of mutations in the γ 2 subunit of AMP-
853 activated protein kinase associated with cardiac hypertrophy and Wolff-Parkinson-
854 White syndrome. *J. Biol. Chem.* 2002;277:51017-51024
- 855 77. Akman HO, Sampayo JN, Ross FA, Scott JW, Wilson G, Benson L, Bruno C, Shanske S,
856 Hardie DG, Dimauro S. Fatal infantile cardiac glycogenosis with phosphorylase kinase
857 deficiency and a mutation in the γ -2 subunit of AMP-activated protein kinase.
858 *Pediatr. Res.* 2007;62:499-504
- 859 78. Burwinkel B, Scott JW, Buhner C, van Landeghem FK, Cox GF, Wilson CJ, Hardie DG,
860 Kilimann MW. Fatal congenital heart glycogenosis caused by a recurrent activating
861 R531Q mutation in the γ 2 subunit of AMP-activated protein kinase (PRKAG2), not by
862 phosphorylase kinase deficiency. *Am. J. Hum. Genet.* 2005;76:1034-1049
- 863 79. Thevenon J, Laurent G, Ader F, Laforet P, Klug D, Duva Pentiah A, Gouya L, Maurage CA,
864 Kacet S, Eicher JC, Albuissou J, Desnos M, Bieth E, Duboc D, Martin L, Reant P, Picard F,
865 Bonithon-Kopp C, Gautier E, Binguet C, Thauvin-Robinet C, Faivre L, Bouvagnet P,
866 Charron P, Richard P. High prevalence of arrhythmic and myocardial complications in
867 patients with cardiac glycogenosis due to PRKAG2 mutations. *Europace.* 2016

- 868 80. Gollob MH, Seger JJ, Gollob TN, Tapscott T, Gonzales O, Bachinski L, Roberts R. Novel
869 PRKAG2 mutation responsible for the genetic syndrome of ventricular preexcitation and
870 conduction system disease with childhood onset and absence of cardiac hypertrophy.
871 *Circulation*. 2001;104:3030-3033
- 872 81. Cheung PCF, Salt IP, Davies SP, Hardie DG, Carling D. Characterization of AMP-activated
873 protein kinase γ subunit isoforms and their role in AMP binding. *Biochem. J.*
874 2000;346:659-669
- 875 82. Luptak I, Shen M, He H, Hirshman MF, Musi N, Goodyear LJ, Yan J, Wakimoto H, Morita H,
876 Arad M, Seidman CE, Seidman JG, Ingwall JS, Balschi JA, Tian R. Aberrant activation of
877 AMP-activated protein kinase remodels metabolic network in favor of cardiac glycogen
878 storage. *J. Clin. Invest.* 2007;117:1432-1439
- 879 83. Kim M, Hunter RW, Garcia-Menendez L, Gong G, Yang YY, Kolwicz SC, Jr., Xu J, Sakamoto
880 K, Wang W, Tian R. Mutation in the γ 2-subunit of AMP-activated protein kinase
881 stimulates cardiomyocyte proliferation and hypertrophy independent of glycogen
882 storage. *Circ. Res.* 2014;114:966-975
- 883 84. Dalle Pezze P, Ruf S, Sonntag AG, Langelaar-Makkinje M, Hall P, Heberle AM, Razquin
884 Navas P, van Eunen K, Tolle RC, Schwarz JJ, Wiese H, Warscheid B, Deitersen J, Stork B,
885 Fassler E, Schauble S, Hahn U, Horvatovich P, Shanley DP, Thedieck K. A systems study
886 reveals concurrent activation of AMPK and mTOR by amino acids. *Nat. Commun.*
887 2016;7:13254
- 888 85. Arad M, Moskowitz IP, Patel VV, Ahmad F, Perez-Atayde AR, Sawyer DB, Walter M, Li GH,
889 Burgon PG, Maguire CT, Stapleton D, Schmitt JP, Guo XX, Pizard A, Kupersmidt S, Roden
890 DM, Berul CI, Seidman CE, Seidman JG. Transgenic mice overexpressing mutant PRKAG2
891 define the cause of Wolff-Parkinson-White syndrome in glycogen storage
892 cardiomyopathy. *Circulation*. 2003;107:2850-2856
- 893 86. Yavari A, Stocker CJ, Ghaffari S, Wargent ET, Steeples V, Czibik G, Pinter K, Bellahcene M,
894 Woods A, Martinez de Morentin PB, Cansell C, Lam BY, Chuster A, Petkevicius K, Nguyen-
895 Tu MS, Martinez-Sanchez A, Pullen TJ, Oliver PL, Stockenhuber A, Nguyen C, Lazdam M,
896 O'Dowd JF, Harikumar P, Toth M, Beall C, Kyriakou T, Parnis J, Sarma D, Katritsis G,
897 Wortmann DD, Harper AR, Brown LA, Willows R, Gandra S, Poncio V, de Oliveira
898 Figueiredo MJ, Qi NR, Peirson SN, McCrimmon RJ, Gereben B, Tretter L, Fekete C,
899 Redwood C, Yeo GS, Heisler LK, Rutter GA, Smith MA, Withers DJ, Carling D, Sternick EB,
900 Arch JR, Cawthorne MA, Watkins H, Ashrafian H. Chronic activation of γ 2 AMPK
901 induces obesity and reduces beta cell function. *Cell Metab.* 2016;23:821-836
- 902 87. Yang X, Mudgett J, Bou-About G, Champy MF, Jacobs H, Monassier L, Pavlovic G, Sorg T,
903 Herault Y, Petit-Demouliere B, Lu K, Feng W, Wang H, Ma LJ, Askew R, Erion MD, Kelley
904 DE, Myers RW, Li C, Guan HP. Physiological expression of AMPK γ 2RG mutation
905 causes Wolff-Parkinson-White syndrome and induces kidney injury in mice. *J. Biol.*
906 *Chem.* 2016;291:23428-23439

- 907 88. Yang Y, Atasoy D, Su HH, Sternson SM. Hunger states switch a flip-flop memory circuit
908 via a synaptic AMPK-dependent positive feedback loop. *Cell*. 2011;146:992-1003
- 909 89. Andersson U, Filipsson K, Abbott CR, Woods A, Smith K, Bloom SR, Carling D, Small CJ.
910 AMP-activated protein kinase plays a role in the control of food intake. *J. Biol. Chem*.
911 2004;279:12005-12008
- 912 90. Milan D, Jeon JT, Looft C, Amarger V, Robic A, Thelander M, Rogel-Gaillard C, Paul S,
913 Iannuccelli N, Rask L, Ronne H, Lundstrom K, Reinsch N, Gellin J, Kalm E, Roy PL,
914 Chardon P, Andersson L. A mutation in PRKAG3 associated with excess glycogen content
915 in pig skeletal muscle. *Science*. 2000;288:1248-1251
- 916 91. Costford SR, Kavaslar N, Ahituv N, Chaudhry SN, Schackwitz WS, Dent R, Pennacchio LA,
917 McPherson R, Harper ME. Gain-of-function R225W mutation in human AMPKgamma3
918 causing increased glycogen and decreased triglyceride in skeletal muscle. *PLoS ONE*.
919 2007;2:e903
- 920 92. Crawford SA, Costford SR, Aguer C, Thomas SC, deKemp RA, DaSilva JN, Lafontaine D,
921 Kendall M, Dent R, Beanlands RS, McPherson R, Harper ME. Naturally occurring R225W
922 mutation of the gene encoding AMP-activated protein kinase (AMPK)gamma(3) results
923 in increased oxidative capacity and glucose uptake in human primary myotubes.
924 *Diabetologia*. 2010;53:1986-1997
- 925 93. Morrow VA, Foufelle F, Connell JM, Petrie JR, Gould GW, Salt IP. Direct activation of AMP-
926 activated protein kinase stimulates nitric-oxide synthesis in human aortic endothelial
927 cells. *J. Biol. Chem*. 2003;278:31629-31639
- 928 94. Stahmann N, Woods A, Spengler K, Heslegrave A, Bauer R, Krause S, Viollet B, Carling D,
929 Heller R. Activation of AMP-activated protein kinase by vascular endothelial growth
930 factor mediates endothelial angiogenesis independently of nitric-oxide synthase. *J. Biol.*
931 *Chem*. 2010;285:10638-10652
- 932 95. Dong Y, Zhang M, Liang B, Xie Z, Zhao Z, Asfa S, Choi HC, Zou MH. Reduction of AMP-
933 activated protein kinase alpha2 increases endoplasmic reticulum stress and
934 atherosclerosis in vivo. *Circulation*. 2010;121:792-803
- 935 96. Bess E, Fisslthaler B, Fromel T, Fleming I. Nitric oxide-induced activation of the AMP-
936 activated protein kinase alpha2 subunit attenuates IkappaB kinase activity and
937 inflammatory responses in endothelial cells. *PLoS One*. 2011;6:e20848
- 938 97. Wang S, Zhang M, Liang B, Xu J, Xie Z, Liu C, Viollet B, Yan D, Zou MH. AMPK{alpha}2
939 deletion causes aberrant expression and activation of NAD(P)H oxidase and consequent
940 endothelial dysfunction in vivo. Role of 26S proteasomes. *Circ. Res*. 2010
- 941 98. Nagata D, Mogi M, Walsh K. AMP-activated protein kinase (AMPK) signaling in
942 endothelial cells is essential for angiogenesis in response to hypoxic stress. *J. Biol. Chem*.
943 2003;278:31000-31006

- 944 99. Wang J, Alexanian A, Ying R, Kizhakekuttu TJ, Dharmashankar K, Vasquez-Vivar J,
945 Gutterman DD, Widlansky ME. Acute exposure to low glucose rapidly induces
946 endothelial dysfunction and mitochondrial oxidative stress: role for AMP kinase.
947 *Arteriosclerosis Thromb. Vasc. Biol.* 2012;32:712-720
- 948 100. Fleming I, Fisslthaler B, Dixit M, Busse R. Role of PECAM-1 in the shear-stress-induced
949 activation of Akt and the endothelial nitric oxide synthase (eNOS) in endothelial cells. *J.*
950 *Cell Sci.* 2005;118:4103-4111
- 951 101. Ouchi N, Kobayashi H, Kihara S, Kumada M, Sato K, Inoue T, Funahashi T, Walsh K.
952 Adiponectin stimulates angiogenesis by promoting cross-talk between AMP-activated
953 protein kinase and Akt signaling in endothelial cells. *J. Biol. Chem.* 2004;279:1304-1309
- 954 102. Day RM, Lee YH, Han L, Kim YC, Feng YH. Angiotensin II activates AMPK for execution of
955 apoptosis through energy-dependent and -independent mechanisms. *Am. J. Physiol. Lung*
956 *Cell. Mol. Physiol.* 2011;301:L772-L781
- 957 103. Xu X, Jhun BS, Ha CH, Jin ZG. Molecular mechanisms of ghrelin-mediated endothelial
958 nitric oxide synthase activation. *Endocrinology.* 2008;149:4183-4192
- 959 104. Stahmann N, Woods A, Carling D, Heller R. Thrombin activates AMP-activated protein
960 kinase in endothelial cells via a pathway involving Ca²⁺/calmodulin-dependent protein
961 kinase kinase beta. *Mol. Cell. Biol.* 2006;26:5933-5945
- 962 105. Levine YC, Li GK, Michel T. Agonist-modulated regulation of AMP-activated protein
963 kinase (AMPK) in endothelial cells. Evidence for an AMPK -> Rac1 -> Akt -> endothelial
964 nitric-oxide synthase pathway. *J. Biol. Chem.* 2007;282:20351-20364
- 965 106. Mount PF, Lane N, Venkatesan S, Steinberg GR, Fraser SA, Kemp BE, Power DA.
966 Bradykinin stimulates endothelial cell fatty acid oxidation by CaMKK-dependent
967 activation of AMPK. *Atherosclerosis.* 2008;200:28-36
- 968 107. Yang S, Wang J. Estrogen activates AMP-activated protein kinase in human endothelial
969 cells via ERbeta/Ca/calmodulin-dependent protein kinase kinase beta pathway. *Cell*
970 *biochemistry and biophysics.* 2015;72:701-707
- 971 108. Reihill JA, Ewart MA, Salt IP. The role of AMP-activated protein kinase in the functional
972 effects of vascular endothelial growth factor-A and -B in human aortic endothelial cells.
973 *Vasc. Cell.* 2011;3:9
- 974 109. Ceolotto G, Gallo A, Papparella I, Franco L, Murphy E, Iori E, Pagnin E, Fadini GP, Albiero
975 M, Semplicini A, Avogaro A. Rosiglitazone reduces glucose-induced oxidative stress
976 mediated by NAD(P)H oxidase via AMPK-dependent mechanism. *Arterioscler. Thromb.*
977 *Vasc. Biol.* 2007;27:2627-2633
- 978 110. Rutherford C, Speirs C, Williams JJ, Ewart MA, Mancini SJ, Hawley SA, Delles C, Viollet B,
979 Costa-Pereira AP, Baillie GS, Salt IP, Palmer TM. Phosphorylation of Janus kinase 1
980 (JAK1) by AMP-activated protein kinase (AMPK) links energy sensing to anti-
981 inflammatory signaling. *Sci. Signal.* 2016;9:ra109

- 982 111. Hwang HJ, Chung HS, Jung TW, Ryu JY, Hong HC, Seo JA, Kim SG, Kim NH, Choi KM, Choi
983 DS, Baik SH, Yoo HJ. The dipeptidyl peptidase-IV inhibitor inhibits the expression of
984 vascular adhesion molecules and inflammatory cytokines in HUVECs via Akt- and AMPK-
985 dependent mechanisms. *Mol. Cell. Endocrinol.* 2015;405:25-34
- 986 112. Krasner NM, Ido Y, Ruderman NB, Cacicedo JM. Glucagon-like peptide-1 (GLP-1) analog
987 liraglutide inhibits endothelial cell inflammation through a calcium and AMPK
988 dependent mechanism. *PLoS One.* 2014;9:e97554
- 989 113. Sun W, Lee TS, Zhu M, Gu C, Wang Y, Zhu Y, Shyy JY. Statins activate AMP-activated
990 protein kinase in vitro and in vivo. *Circulation.* 2006;114:2655-2662
- 991 114. Murakami H, Murakami R, Kambe F, Cao X, Takahashi R, Asai T, Hirai T, Numaguchi Y,
992 Okumura K, Seo H, Murohara T. Fenofibrate activates AMPK and increases eNOS
993 phosphorylation in HUVEC. *Biochem. Biophys. Res. Commun.* 2006;341:973-978
- 994 115. Wu Y, Song P, Xu J, Zhang M, Zou MH. Activation of protein phosphatase 2A by palmitate
995 inhibits AMP-activated protein kinase. *J. Biol. Chem.* 2007;282:9777-9788
- 996 116. Weikel KA, Cacicedo JM, Ruderman NB, Ido Y. Glucose and palmitate uncouple AMPK
997 from autophagy in human aortic endothelial cells. *Am. J. Physiol. Cell Physiol.*
998 2015;308:C249-C263
- 999 117. Cacicedo JM, Yagihashi N, Keaney JF, Jr., Ruderman NB, Ido Y. AMPK inhibits fatty acid-
1000 induced increases in NF-kappaB transactivation in cultured human umbilical vein
1001 endothelial cells. *Biochem. Biophys. Res. Commun.* 2004;324:1204-1209
- 1002 118. Heathcote HR, Mancini SJ, Strembitska A, Jamal K, Reihill JA, Palmer TM, Gould GW, Salt
1003 IP. Protein kinase C phosphorylates AMP-activated protein kinase alpha1 Ser487.
1004 *Biochem. J.* 2016;473:4681-4697
- 1005 119. Dagher Z, Ruderman N, Tornheim K, Ido Y. The effect of AMP-activated protein kinase
1006 and its activator AICAR on the metabolism of human umbilical vein endothelial cells.
1007 *Biochem. Biophys. Res. Commun.* 1999;265:112-115
- 1008 120. Ido Y, Carling D, Ruderman N. Hyperglycemia-induced apoptosis in human umbilical
1009 vein endothelial cells: inhibition by the AMP-activated protein kinase activation.
1010 *Diabetes.* 2002;51:159-167
- 1011 121. Goirand F, Solar M, Athea Y, Viollet B, Mateo P, Fortin D, Leclerc J, Hoerter J, Ventura-
1012 Clapier R, Garnier A. Activation of AMP kinase alpha1 subunit induces aortic
1013 vasorelaxation in mice. *J. Physiol.* 2007;581:1163-1171
- 1014 122. Song P, Wang S, He C, Liang B, Viollet B, Zou MH. AMPK-alpha2 deletion exacerbates
1015 neointima formation by upregulating Skp2 in vascular smooth muscle cells. *Circ. Res.*
1016 2011;109:1230-1239
- 1017 123. Motobayashi Y, Izawa-Ishizawa Y, Ishizawa K, Orino S, Yamaguchi K, Kawazoe K,
1018 Hamano S, Tsuchiya K, Tomita S, Tamaki T. Adiponectin inhibits insulin-like growth
1019 factor-1-induced cell migration by the suppression of extracellular signal-regulated
1020 kinase 1/2 activation, but not Akt in vascular smooth muscle cells. *Hypertens. Res.*
1021 2009;32:188-193

- 1022 124. Ning J, Clemmons DR. AMP-activated protein kinase inhibits IGF-I signaling and protein
1023 synthesis in vascular smooth muscle cells via stimulation of insulin receptor substrate 1
1024 S794 and tuberous sclerosis 2 S1345 phosphorylation. *Mol. Endocrinol.* 2010;24:1218-
1025 1229
- 1026 125. Gayard M, Guilluy C, Rousselle A, Viollet B, Henrion D, Pacaud P, Loirand G, Rolli-
1027 Derkinderen M. AMPK alpha 1-induced RhoA phosphorylation mediates vasoprotective
1028 effect of estradiol. *Arterioscler. Thromb. Vasc. Biol.* 2011;31:2634-2642
- 1029 126. Pyla R, Pichavaram P, Fairaq A, Park MA, Kozak M, Kamath V, Patel VS, Segar L. Altered
1030 energy state reversibly controls smooth muscle contractile function in human
1031 saphenous vein during acute hypoxia-reoxygenation: Role of glycogen, AMP-activated
1032 protein kinase, and insulin-independent glucose uptake. *Biochem. Pharmacol.*
1033 2015;97:77-88
- 1034 127. Nagata D, Takeda R, Sata M, Satonaka H, Suzuki E, Nagano T, Hirata Y. AMP-activated
1035 protein kinase inhibits angiotensin II-stimulated vascular smooth muscle cell
1036 proliferation. *Circulation.* 2004;110:444-451
- 1037 128. Kim JE, Choi HC. Losartan Inhibits vascular smooth muscle cell proliferation through
1038 activation of AMP-activated protein kinase. *The Korean journal of physiology &*
1039 *pharmacology : official journal of the Korean Physiological Society and the Korean Society*
1040 *of Pharmacology.* 2010;14:299-304
- 1041 129. Ning J, Xi G, Clemmons DR. Suppression of AMPK activation via S485 phosphorylation by
1042 IGF-I during hyperglycemia is mediated by AKT activation in vascular smooth muscle
1043 cells. *Endocrinology.* 2011;152:3143-3154
- 1044 130. Son BK, Akishita M, Iijima K, Kozaki K, Maemura K, Eto M, Ouchi Y. Adiponectin
1045 antagonizes stimulatory effect of tumor necrosis factor-alpha on vascular smooth
1046 muscle cell calcification: regulation of growth arrest-specific gene 6-mediated survival
1047 pathway by adenosine 5'-monophosphate-activated protein kinase. *Endocrinology.*
1048 2008;149:1646-1653
- 1049 131. Cao X, Li H, Tao H, Wu N, Yu L, Zhang D, Lu X, Zhu J, Lu Z, Zhu Q. Metformin inhibits
1050 vascular calcification in female rat aortic smooth muscle cells via the AMPK-eNOS-NO
1051 pathway. *Endocrinology.* 2013;154:3680-3689
- 1052 132. Siragusa M, Fleming I. The eNOS signalosome and its link to endothelial dysfunction.
1053 *Pflugers Arch.* 2016;468:1125-1137
- 1054 133. Chen ZP, Mitchelhill KI, Michell BJ, Stapleton D, Rodriguez-Crespo I, Witters LA, Power
1055 DA, Ortiz de Montellano PR, Kemp BE. AMP-activated protein kinase phosphorylation of
1056 endothelial NO synthase. *FEBS Lett.* 1999;443:285-289
- 1057 134. Chen Z, Peng IC, Sun W, Su MI, Hsu PH, Fu Y, Zhu Y, Defea K, Pan S, Tsai MD, Shyy JY.
1058 AMP-activated protein kinase functionally phosphorylates endothelial nitric oxide
1059 synthase Ser633. *Circ. Res.* 2009;104:496-505

- 1060 135. Wang S, Xu J, Song P, Viollet B, Zou MH. In vivo activation of AMP-activated protein
1061 kinase attenuates diabetes-enhanced degradation of GTP cyclohydrolase I. *Diabetes*.
1062 2009;58:1893-1901
- 1063 136. Mount PF, Hill RE, Fraser SA, Levidiotis V, Katsis F, Kemp BE, Power DA. Acute renal
1064 ischemia rapidly activates the energy sensor AMPK but does not increase
1065 phosphorylation of eNOS-Ser1177. *Am. J. Physiol. Renal Physiol.* 2005;289:F1103-F1115
- 1066 137. Kukidome D, Nishikawa T, Sonoda K, Imoto K, Fujisawa K, Yano M, Motoshima H,
1067 Taguchi T, Matsumura T, Araki E. Activation of AMP-activated protein kinase reduces
1068 hyperglycemia-induced mitochondrial reactive oxygen species production and promotes
1069 mitochondrial biogenesis in human umbilical vein endothelial cells. *Diabetes*.
1070 2006;55:120-127
- 1071 138. Shah MS, Brownlee M. Molecular and cellular mechanisms of cardiovascular disorders in
1072 diabetes. *Circ. Res.* 2016;118:1808-1829
- 1073 139. Colombo SL, Moncada S. AMPK α 1 regulates the antioxidant status of vascular
1074 endothelial cells. *Biochem. J.* 2009;421:163-169
- 1075 140. Hou X, Song J, Li XN, Zhang L, Wang X, Chen L, Shen YH. Metformin reduces intracellular
1076 reactive oxygen species levels by upregulating expression of the antioxidant thioredoxin
1077 via the AMPK-FOXO3 pathway. *Biochem. Biophys. Res. Commun.* 2010;396:199-205
- 1078 141. Dong Y, Zhang M, Wang S, Liang B, Zhao Z, Liu C, Wu M, Choi HC, Lyons TJ, Zou MH.
1079 Activation of AMP-activated protein kinase inhibits oxidized LDL-triggered endoplasmic
1080 reticulum stress in vivo. *Diabetes*. 2010;59:1386-1396
- 1081 142. Ridker PM, Luscher TF. Anti-inflammatory therapies for cardiovascular disease. *Eur.*
1082 *Heart J.* 2014;35:1782-1791
- 1083 143. Salt IP, Palmer TM. Exploiting the anti-inflammatory effects of AMP-activated protein
1084 kinase activation. *Expert Opin. Investig. Drugs.* 2012;21:1155-1167
- 1085 144. Ewart MA, Kohlhaas CF, Salt IP. Inhibition of tumor necrosis factor α -stimulated
1086 monocyte adhesion to human aortic endothelial cells by AMP-activated protein kinase.
1087 *Arterioscler. Thromb. Vasc. Biol.* 2008;28:2255-2257
- 1088 145. Spiecker M, Peng HB, Liao JK. Inhibition of endothelial vascular cell adhesion molecule-1
1089 expression by nitric oxide involves the induction and nuclear translocation of
1090 IkappaB α . *J. Biol. Chem.* 1997;272:30969-30974
- 1091 146. Zhang Y, Qiu J, Wang X, Xia M. AMP-activated protein kinase suppresses endothelial cell
1092 inflammation through phosphorylation of transcriptional coactivator p300. *Arterioscler.*
1093 *Thromb. Vasc. Biol.* 2011;31:2897-2908
- 1094 147. Schulz E, Dopheide J, Schuhmacher S, Thomas SR, Chen K, Daiber A, Wenzel P, Munzel T,
1095 Keaney JF, Jr. Suppression of the JNK pathway by induction of a metabolic stress
1096 response prevents vascular injury and dysfunction. *Circulation.* 2008;118:1347-1357

- 1097 148. Mancini SJ, White AD, Bijland S, Rutherford C, Graham D, Richter EA, Viollet B, Touyz RM,
1098 Palmer TM, Salt IP. Activation of AMP-activated protein kinase rapidly suppresses
1099 multiple pro-inflammatory pathways in adipocytes including IL-1 receptor-associated
1100 kinase-4 phosphorylation. *Mol. Cell. Endocrinol.* 2017;440:44-56
- 1101 149. Kim SA, Choi HC. Metformin inhibits inflammatory response via AMPK-PTEN pathway in
1102 vascular smooth muscle cells. *Biochem. Biophys. Res. Commun.* 2012;425:866-872
- 1103 150. He C, Li H, Viollet B, Zou MH, Xie Z. AMPK suppresses vascular inflammation in vivo by
1104 inhibiting Signal Transducer and Activator of Transcription-1. *Diabetes.* 2015;64:4285-
1105 4297
- 1106 151. Zippel N, Malik RA, Fromel T, Popp R, Bess E, Strilic B, Wettschureck N, Fleming I,
1107 Fisslthaler B. Transforming growth factor-beta-activated kinase 1 regulates
1108 angiogenesis via AMP-activated protein kinase-alpha1 and redox balance in endothelial
1109 cells. *Arterioscler. Thromb. Vasc. Biol.* 2013;33:2792-2799
- 1110 152. Xu MJ, Song P, Shirwany N, Liang B, Xing J, Viollet B, Wang X, Zhu Y, Zou MH. Impaired
1111 expression of uncoupling protein 2 causes defective postischemic angiogenesis in mice
1112 deficient in AMP-activated protein kinase alpha subunits. *Arterioscler. Thromb. Vasc.*
1113 *Biol.* 2011;31:1757-1765
- 1114 153. Ouchi N, Shibata R, Walsh K. AMP-activated protein kinase signaling stimulates VEGF
1115 expression and angiogenesis in skeletal muscle. *Circ. Res.* 2005;96:838-846
- 1116 154. Wang S, Liang B, Viollet B, Zou MH. Inhibition of the AMP-activated protein kinase-
1117 alpha2 accentuates agonist-induced vascular smooth muscle contraction and high blood
1118 pressure in mice. *Hypertension.* 2011;57:1010-1017
- 1119 155. Schneider H, Schubert KM, Blodow S, Kreutz CP, Erdogmus S, Wiedenmann M, Qiu J, Fey
1120 T, Ruth P, Lubomirov LT, Pfitzer G, Mederos Y, M. S, Hardie DG, Gudermann T, Pohl U.
1121 AMPK dilates resistance arteries via activation of SERCA and BKCa channels in smooth
1122 muscle. *Hypertension.* 2015;66:108-116
- 1123 156. Igata M, Motoshima H, Tsuruzoe K, Kojima K, Matsumura T, Kondo T, Taguchi T,
1124 Nakamaru K, Yano M, Kukidome D, Matsumoto K, Toyonaga T, Asano T, Nishikawa T,
1125 Araki E. Adenosine monophosphate-activated protein kinase suppresses vascular
1126 smooth muscle cell proliferation through the inhibition of cell cycle progression. *Circ.*
1127 *Res.* 2005;97:837-844
- 1128 157. Song P, Wang S, He C, Liang B, Viollet B, Zou MH. AMPKalpha2 deletion exacerbates
1129 neointima formation by upregulating Skp2 in vascular smooth muscle cells. *Circ. Res.*
1130 2011;109:1230-1239
- 1131 158. Ding Y, Zhang M, Zhang W, Lu Q, Cai Z, Song P, Okon IS, Xiao L, Zou MH. AMP-activated
1132 protein kinase alpha 2 deletion induces VSMC phenotypic switching and reduces
1133 features of atherosclerotic plaque stability. *Circ. Res.* 2016;119:718-730

- 1134 159. Song P, Zhou Y, Coughlan KA, Dai X, Xu H, Viollet B, Zou MH. Adenosine monophosphate-
1135 activated protein kinase- α 2 deficiency promotes vascular smooth muscle cell
1136 migration via S-phase kinase-associated protein 2 upregulation and E-cadherin
1137 downregulation. *Arterioscler. Thromb. Vasc. Biol.* 2013;33:2800-2809
- 1138 160. Cai Z, Ding Y, Zhang M, Lu Q, Wu S, Zhu H, Song P, Zou MH. Ablation of adenosine
1139 monophosphate-activated protein kinase α 1 in vascular smooth muscle cells
1140 promotes diet-Induced atherosclerotic calcification in vivo. *Circ. Res.* 2016;119:422-433
- 1141 161. Cacicedo JM, Gauthier MS, Lebrasseur NK, Jasuja R, Ruderman NB, Ido Y. Acute exercise
1142 activates AMPK and eNOS in the mouse aorta. *Am. J. Physiol. Heart Circ. Physiol.*
1143 2011;301:H1255-H1265
- 1144 162. Ma L, Ma S, He H, Yang D, Chen X, Luo Z, Liu D, Zhu Z. Perivascular fat-mediated vascular
1145 dysfunction and remodeling through the AMPK/mTOR pathway in high-fat diet-induced
1146 obese rats. *Hypertens. Res.*
- 1147 163. Zhao CX, Xu X, Cui Y, Wang P, Wei X, Yang S, Edin ML, Zeldin DC, Wang DW. Increased
1148 endothelial nitric-oxide synthase expression reduces hypertension and
1149 hyperinsulinemia in fructose-treated rats. *J. Pharmacol. Exp. Ther.* 2009;328:610-620
- 1150 164. Bradley EA, Eringa EC, Stehouwer CD, Korstjens I, van Nieuw Amerongen GP, Musters R,
1151 Sipkema P, Clark MG, Rattigan S. Activation of AMP-activated protein kinase by 5-
1152 aminoimidazole-4-carboxamide-1- β -D-ribofuranoside in the muscle microcirculation
1153 increases nitric oxide synthesis and microvascular perfusion. *Arterioscler. Thromb. Vasc.*
1154 *Biol.* 2010;30:1137-1142
- 1155 165. Ford RJ, Teschke SR, Reid EB, Durham KK, Kroetsch JT, Rush JW. AMP-activated protein
1156 kinase activator AICAR acutely lowers blood pressure and relaxes isolated resistance
1157 arteries of hypertensive rats. *J. Hypertens.* 2012;30:725-733
- 1158 166. Ford RJ, Rush JW. Endothelium-dependent vasorelaxation to the AMPK activator AICAR
1159 is enhanced in aorta from hypertensive rats and is NO and EDCF dependent. *Am. J.*
1160 *Physiol. Heart Circ. Physiol.* 2011;300:H64-H75
- 1161 167. Bosselaar M, Boon H, van Loon LJ, van den Broek PH, Smits P, Tack CJ. Intra-arterial
1162 AICA-riboside administration induces NO-dependent vasodilation in vivo in human
1163 skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* 2009;297:E759-E766
- 1164 168. Enkhjargal B, Godo S, Sawada A, Suvd N, Saito H, Noda K, Satoh K, Shimokawa H.
1165 Endothelial AMP-activated protein kinase regulates blood pressure and coronary flow
1166 responses through hyperpolarization mechanism in mice. *Arterioscler. Thromb. Vasc.*
1167 *Biol.* 2014;34:1505-1513
- 1168 169. Liang B, Wang S, Wang Q, Zhang W, Viollet B, Zhu Y, Zou MH. Aberrant endoplasmic
1169 reticulum stress in vascular smooth muscle increases vascular contractility and blood
1170 pressure in mice deficient of AMP-activated protein kinase- α 2 in vivo. *Arterioscler.*
1171 *Thromb. Vasc. Biol.* 2013;33:595-604

- 1172 170. Kohlstedt K, Trouvain C, Boettger T, Shi L, Fisslthaler B, Fleming I. AMP-activated
1173 protein kinase regulates endothelial cell angiotensin-converting enzyme expression via
1174 p53 and the post-transcriptional regulation of microRNA-143/145. *Circ. Res.*
1175 2013;112:1150-1158
- 1176 171. Buhl ES, Jessen N, Pold R, Ledet T, Flyvbjerg A, Pedersen SB, Pedersen O, Schmitz O,
1177 Lund S. Long-term AICAR administration reduces metabolic disturbances and lowers
1178 blood pressure in rats displaying features of the insulin resistance syndrome. *Diabetes.*
1179 2002;51:2199-2206
- 1180 172. Meijer RI, Bakker W, Alta CL, Sipkema P, Yudkin JS, Viollet B, Richter EA, Smulders YM,
1181 van Hinsbergh VW, Serne EH, Eringa EC. Perivascular adipose tissue control of insulin-
1182 induced vasoreactivity in muscle is impaired in db/db mice. *Diabetes.* 2013;62:590-598
- 1183 173. Almagrouk TA, Ugusman AB, Katwan OJ, Salt IP, Kennedy S. Deletion of AMPK α 1
1184 attenuates the anticontractile effect of perivascular adipose tissue (PVAT) and reduces
1185 adiponectin release. *Br. J. Pharmacol.* 2017;in press
- 1186 174. Gaskin FS, Kamada K, Zuidema MY, Jones AW, Rubin LJ, Korthuis RJ. Isoform-selective 5'-
1187 AMP-activated protein kinase-dependent preconditioning mechanisms to prevent
1188 postischemic leukocyte-endothelial cell adhesive interactions. *Am. J. Physiol. Heart Circ.*
1189 *Physiol.* 2011;300:H1352-H1360
- 1190 175. Li D, Wang D, Wang Y, Ling W, Feng X, Xia M. Adenosine monophosphate-activated
1191 protein kinase induces cholesterol efflux from macrophage-derived foam cells and
1192 alleviates atherosclerosis in apolipoprotein E-deficient mice. *J. Biol. Chem.*
1193 2010;285:33499-33509
- 1194 176. Chen B, Li J, Zhu H. AMP-activated protein kinase attenuates oxLDL uptake in
1195 macrophages through PP2A/NF-kappaB/LOX-1 pathway. *Vascul. Pharmacol.* 2016;85:1-
1196 10
- 1197 177. Vasamsetti SB, Karnewar S, Kanugula AK, Thatipalli AR, Kumar JM, Kotamraju S.
1198 Metformin inhibits monocyte-to-macrophage differentiation via AMPK-mediated
1199 inhibition of STAT3 Activation: potential role in atherosclerosis. *Diabetes.*
1200 2015;64:2028-2041
- 1201 178. Wang Q, Zhang M, Liang B, Shirwany N, Zhu Y, Zou MH. Activation of AMP-activated
1202 protein kinase is required for berberine-induced reduction of atherosclerosis in mice:
1203 the role of uncoupling protein 2. *PLoS One.* 2011;6:e25436
- 1204 179. UKPDS. Effect of intensive blood-glucose control with metformin on complications in
1205 overweight patients with type 2 diabetes (UKPDS 34). UK Prospective Diabetes Study
1206 (UKPDS) Group. *Lancet.* 1998;352:854-865
- 1207 180. Dai X, Ding Y, Liu Z, Zhang W, Zou MH. Phosphorylation of CHOP (C/EBP Homologous
1208 Protein) by the AMP-activated protein kinase α 1 in macrophages promotes CHOP
1209 degradation and reduces injury-induced neointimal disruption in vivo. *Circ. Res.*
1210 2016;119:1089-1100

- 1211 181. Li FY, Lam KS, Tse HF, Chen C, Wang Y, Vanhoutte PM, Xu A. Endothelium-selective
1212 activation of AMP-activated protein kinase prevents diabetes mellitus-induced
1213 impairment in vascular function and reendothelialization via induction of heme
1214 oxygenase-1 in mice. *Circulation*. 2012;126:1267-1277
- 1215 182. Cao Q, Cui X, Wu R, Zha L, Wang X, Parks JS, Yu L, Shi H, Xue B. Myeloid deletion of
1216 alpha1AMPK exacerbates atherosclerosis in LDL receptor knockout (LDLRKO) mice.
1217 *Diabetes*. 2016;65:1565-1576
- 1218 183. Song Y, Wu Y, Su X, Zhu Y, Liu L, Pan Y, Zhu B, Yang L, Gao L, Li M. Activation of AMPK
1219 inhibits PDGF-induced pulmonary arterial smooth muscle cells proliferation and its
1220 potential mechanisms. *Pharmacol. Res.* 2016;107:117-124
- 1221 184. Dean A, Nilsen M, Loughlin L, Salt IP, MacLean MR. Metformin reverses development of
1222 pulmonary hypertension via aromatase inhibition. *Hypertension*. 2016;68:446-454
- 1223 185. Omura J, Satoh K, Kikuchi N, Satoh T, Kurosawa R, Nogi M, Otsuki T, Kozu K, Numano K,
1224 Suzuki K, Sunamura S, Tatebe S, Aoki T, Sugimura K, Miyata S, Hoshikawa Y, Okada Y,
1225 Shimokawa H. Protective roles of endothelial AMP-activated protein kinase against
1226 hypoxia-induced pulmonary hypertension in mice. *Circ. Res.* 2016;119:197-209
- 1227 186. Ibe JC, Zhou Q, Chen T, Tang H, Yuan JX, Raj JU, Zhou G. Adenosine monophosphate-
1228 activated protein kinase is required for pulmonary artery smooth muscle cell survival
1229 and the development of hypoxic pulmonary hypertension. *Am. J. Respir. Cell. Mol. Biol.*
1230 2013;49:609-618
- 1231 187. Moral-Sanz J, Mahmoud AD, Ross FA, Eldstrom J, Fedida D, Hardie DG, Evans AM. AMP-
1232 activated protein kinase inhibits Kv 1.5 channel currents of pulmonary arterial myocytes
1233 in response to hypoxia and inhibition of mitochondrial oxidative phosphorylation. *J.*
1234 *Physiol.* 2016;594:4901-4915
- 1235 188. Pinter K, Grignani RT, Czibik G, Farza H, Watkins H, Redwood C. Embryonic expression
1236 of AMPK gamma subunits and the identification of a novel gamma2 transcript variant in
1237 adult heart. *J. Mol. Cell. Cardiol.* 2012;53:342-349
- 1238 189. Cool B, Zinker B, Chiou W, Kifle L, Cao N, Perham M, Dickinson R, Adler A, Gagne G,
1239 Iyengar R, Zhao G, Marsh K, Kym P, Jung P, Camp HS, Frevert E. Identification and
1240 characterization of a small molecule AMPK activator that treats key components of type
1241 2 diabetes and the metabolic syndrome. *Cell Metab.* 2006;3:403-416
- 1242 190. Zadra G, Photopoulos C, Tyekucheva S, Heidari P, Weng QP, Fedele G, Liu H, Scaglia N,
1243 Priolo C, Sicinska E, Mahmood U, Signoretti S, Birnberg N, Loda M. A novel direct
1244 activator of AMPK inhibits prostate cancer growth by blocking lipogenesis. *EMBO Mol.*
1245 *Med.* 2014;6:519-538

- 1246 191. Salatto CT, Miller RA, Cameron KO, Cokorinos E, Reyes A, Ward J, Calabrese M,
1247 Kurumbail R, Rajamohan F, Kalgutkar AS, Tess DA, Shavnya A, Genung NE, Edmonds DJ,
1248 Jatkar A, Maciejewski BS, Amaro M, Gandhok H, Monetti M, Cialdea K, Bollinger E,
1249 Kreeger JM, Coskran TM, Opsahl AC, Boucher GG, Birnbaum MJ, DaSilva-Jardine P, Rolph
1250 T. Selective Activation of AMPK b1-containing Isoforms Improves Kidney Function in a
1251 Rat Model of Diabetic Nephropathy. *J. Pharmacol. Exp. Ther.* 2017
- 1252 192. Gollob MH, Green MS, Tang AS, Gollob T, Karibe A, Ali Hassan AS, Ahmad F, Lozado R,
1253 Shah G, Fananapazir L, Bachinski LL, Roberts R. Identification of a gene responsible for
1254 familial Wolff-Parkinson-White syndrome. *N. Engl. J. Med.* 2001;344:1823-1831
- 1255 193. Arad M, D.W. B, Perez-Atayde AR, McKenna WJ, Sparks EA, Kanter RJ, McGarry K,
1256 Seidman JG, Seidman CE. Constitutively active AMP kinase mutations cause glycogen
1257 storage disease mimicking hypertrophic cardiomyopathy. *J. Clin. Invest.* 2002;109:357-
1258 362
- 1259 194. Charron P, Genest M, Richard P, Komajda M, Pochmalicki G. A familial form of
1260 conduction defect related to a mutation in the PRKAG2 gene. *Europace.* 2007;9:597-600
- 1261 195. Blair E, Redwood C, Ashrafian H, Oliveira M, Broxholme J, Kerr B, Salmon A, Ostman-
1262 Smith I, Watkins H. Mutations in the gamma-2 subunit of AMP-activated protein kinase
1263 cause familial hypertrophic cardiomyopathy: evidence for the central role of energy
1264 compromise in disease pathogenesis. *Hum. Mol. Genet.* 2001;10:1215-1220
- 1265 196. Arad M, Maron BJ, Gorham JM, Johnson WH, Jr., Saul JP, Perez-Atayde AR, Spirito P,
1266 Wright GB, Kanter RJ, Seidman CE, Seidman JG. Glycogen storage diseases presenting as
1267 hypertrophic cardiomyopathy. *N. Engl. J. Med.* 2005;352:362-372
- 1268 197. Bayrak F, Komurcu-Bayrak E, Mutlu B, Kahveci G, Basaran Y, Erginel-Unaltuna N.
1269 Ventricular pre-excitation and cardiac hypertrophy mimicking hypertrophic
1270 cardiomyopathy in a Turkish family with a novel PRKAG2 mutation. *Eur. J. Heart Fail.*
1271 2006;8:712-715
- 1272 198. Xie C, Zhang YP, Song L, Luo J, Qi W, Hu J, Lu D, Yang Z, Zhang J, Xiao J, Zhou B, Du JL, Jing
1273 N, Liu Y, Wang Y, Li BL, Song BL, Yan Y. Genome editing with CRISPR/Cas9 in postnatal
1274 mice corrects PRKAG2 cardiac syndrome. *Cell Res.* 2016;26:1099-1111
- 1275 199. Laforet P, Richard P, Said MA, Romero NB, Lacene E, Leroy JP, Baussan C, Hogrel JY,
1276 Lavergne T, Wahbi K, Hainque B, Duboc D. A new mutation in PRKAG2 gene causing
1277 hypertrophic cardiomyopathy with conduction system disease and muscular
1278 glycogenosis. *Neuromuscul. Disord.* 2006;16:178-182
- 1279

1280 **FIGURE LEGENDS**

1281 **Figure 1: Domain layouts of AMPK subunits and their isoforms.** The linear layout of domains
1282 is shown, approximately to scale and with similar color coding as in Fig. 2. Note that
1283 both β subunits are N-myristoylated, and that the $\gamma 2$ and $\gamma 3$ subunit isoforms have
1284 unrelated N-terminal extensions of unknown function, although both are also reported
1285 to exist as shorter, N-terminally truncated versions due to alternate start sites and/or
1286 splicing¹⁸⁸.

1287 **Figure 2: Structure of human AMPK ($\alpha 1\beta 2\gamma 1$ complex)⁷.** The model was created in
1288 spacefilling mode using PyMol v1.7.4.2 with the co-ordinates in PDB file 4RER, and
1289 with color coding similar to Fig. 1. The heterotrimer was crystallized in the presence of
1290 β -cyclodextrin, which occupies the glycogen-binding site, staurosporine, which
1291 occupies the active site, and AMP, which occupies sites 1, 3 and 4 on the γ subunit
1292 (sites 1 and 4 are round the back in this view). Although Thr172 was phosphorylated, it
1293 is not visible in this view but lies in the cleft between the α subunit C lobe and the β -
1294 CTD, just over the right-hand “shoulder” of the C-lobe.

1295 **Figure 3: Two views of the structure of the four CBS repeats of the $\gamma 1$ subunit.** The model
1296 used the co-ordinates in PDB file 2V8Q¹⁸, and was rendered in PyMol v1.7.4.2 with the
1297 $\gamma 1$ subunit in cartoon view. The two views are rotated 180° around the x axis (dashed
1298 line) with respect to each other, with the orientation of the top view being similar to that
1299 in Fig. 2. Note the pseudosymmetrical layout of the four CBS repeats, which are
1300 colored differently (and differently to Figs. 1 and 2). Residues equivalent to those
1301 mutated in $\gamma 2$ are highlighted using the “dots” version of space-filling representation,
1302 and are numbered using the human $\gamma 2$ numbering. The three molecules of bound AMP
1303 are labelled and are shown in standard space-filling view, with C atoms green, O red
1304 and N blue.

1305 **Figure 4: Regulation of endothelial NO and superoxide synthesis by AMPK.** AMPK
1306 activation stimulates NO synthesis via multiple mechanisms: (i) phosphorylation of
1307 eNOS at Ser633 and Ser1177; (ii) increasing Hsp90 association with eNOS; (iii)
1308 increasing BH₄ concentrations via GTP cyclohydrolase I (GTPCH1); (iv) reducing
1309 superoxide synthesis via inhibition of Nox and increasing antioxidant protein

(superoxide dismutase (SOD), thioredoxin and catalase) levels. NO itself is also reported to activate AMPK.

Figure 5: Actions of AMPK in vascular cells. Physiological activators of AMPK in ECs include hypoxia/ischemia, shear stress, adiponectin, thrombin, bradykinin and VEGF (synthesis of which is stimulated by AMPK in other tissues). AMPK is reported to stimulate VSMC relaxation through: (i) increased NO synthesis and possibly endothelium-dependent hypopolarising factor (EDHF); (ii) inhibition of MYPT1/MLC phosphorylation and Ca^{2+} levels in VSMCs, reported to be mediated by reduced RhoA activity and increased sarco/endoplasmic Ca^{2+} ATPase (SERCA) activity respectively. AMPK activation in ECs stimulates proliferation and migration, whereas in VSMCs, proliferation and migration are inhibited, associated with p53 phosphorylation, Rb dephosphorylation and p27(Kip1) stabilisation. AMPK also inhibits VSMC calcification by reducing Runx2 and pro-inflammatory signalling pathways leading to leukocyte adhesion and cytokine/chemokine synthesis.


Table 1: List of pharmacological agents commonly used to activate AMPK in intact cells or in vivo, and their mechanisms of action.


| Class | Agent | Mechanism | Binding site used | Isoform-selective? | Ref. |
|-------|----------------|--------------------------------|-----------------------|--------------------|--------|
| 1 | metformin | mitochondrial inhibitor, AMP ↑ | γ subunit (binds AMP) | No | 33 |
| 1 | phenformin | mitochondrial inhibitor, AMP ↑ | γ subunit (binds AMP) | No | 33 |
| 1 | berberine | mitochondrial inhibitor, AMP ↑ | γ subunit (binds AMP) | No | 33 |
| 2 | 2-deoxyglucose | glycolytic inhibitor, AMP ↑ | γ subunit (binds AMP) | No | 33 |
| 3 | AICAR | pro-drug, converted to ZMP | γ subunit (binds ZMP) | No | 33, 34 |
| 4 | C13 | pro-drug, converted to C2 | γ subunit (binds C2) | α1-selective | 38, 40 |
| 5 | A769662 | direct activator | ADaM site | β1-selective | 189 |
| 5 | 991 | direct activator | ADaM site | β1>β2 | 8 |
| 5 | MT 63-78 | direct activator | ADaM site | β1-selective | 190 |
| 5 | PF-06409577 | direct activator | ADaM site | β1-selective | 191 |
| 5 | PF-249 | direct activator | ADaM site | β1-selective | 191 |
| 5 | salicylate | direct activator | ADaM site | β1-selective | 9, 44 |



Table 2: Amino acid replacements, generated by mutations in *PRKAG2*, that are associated with heart disorders. Equivalent residues in the sequences of human $\gamma 1$ and $\gamma 3$ are shown in the second and third column and the CBS repeat affected in the fourth. V336 and R350 are located in the linker between CBS1 and CBS2.

| $\gamma 2$ mutation | $\gamma 1$ equivalent | $\gamma 3$ equivalent | CBS repeat affected | Reference |
|-----------------------|-----------------------|-----------------------|---------------------|-----------|
| R302Q | R70 | R225 | CBS1 | 192-194 |
| S333P | S101 | S333 | CBS1 | 79 |
| V336A | V104 | V259 | linker | 79 |
| L insert (after R350) | after R118 | after R273 | linker | 195 |
| H383R | H151 | H306 | CBS2 | 195 |
| R384T | R152 | R307 | CBS2 | 77 |
| T400N | T168 | T323 | CBS2 | 193 |
| Y487H | Y255 | Y410 | CBS3 | 196 |
| N488I | N256 | N411 | CBS3 | 193 |
| E506K | E274 | E429 | CBS4 | 79, 197 |
| H530R | H298 | H453 | CBS4 | 198 |
| R531G | R299 | R454 | CBS4 | 80 |
| R531Q | R299 | R454 | CBS4 | 78 |
| S548P | S316 | S471 | CBS4 | 199 |


Salt & Hardie Fig. 1







Salt & Hardie Fig. 4



Salt & Hardie Fig. 5

